

**PCT**

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/12, C07K 14/705, C12N 1/21, 15/70, G01N 33/53, 33/68, A61K 38/17, C07K 17/00</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/32767</b>  <b>(43) International Publication Date:</b> 8 June 2000 (08.06.00)
<b>(21) International Application Number:</b> PCT/EP99/09440  <b>(22) International Filing Date:</b> 3 December 1999 (03.12.99)  <b>(30) Priority Data:</b> 98122969.3 3 December 1998 (03.12.98) EP  <b>(71) Applicant (for all designated States except US):</b> MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN E.V. [DE/DE]; Hofgartenstrasse 8, D-80539 München (DE).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> SONDERMANN, Peter [DE/DE]; Pfeuferstrasse 97, D-81373 München (DE). HUBER, Robert [DE/DE]; Schlesierstrasse 13, D-82110 Germering (DE). JAKOB, Uwe [DE/DE]; Guldeinstrasse 42, D-80339 München (DE).  <b>(74) Agents:</b> WEICKMANN, H. et al.; Kopernikusstrasse 9, D-81679 München (DE).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.          Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> RECOMBINANT SOLUBLE Fc RECEPTORS			
<b>(57) Abstract</b>  Recombinant soluble Fc receptors according to the present invention are characterized by the absence of transmembrane domains, signal peptides and glycosylation. Such Fc receptors can easily be obtained by expressing respective nucleic acids in prokaryotic host cells and renaturation of the obtained inclusion bodies, which procedure leads to a very homogenous and pure product. The products can be used for diagnostic as well as pharmaceutical applications and also for the generation of crystal structure data. Such crystal structure data can be used for the modelling of artificial molecules. A further embodiment comprises coupling the Fc receptors according to the invention to solid materials like chromatography materials that can be used to separate and/or enrich antibodies.			
Application No.: 10/756,153 Attorney Docket No.: 13783-105015 References B9			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**Recombinant soluble Fc receptors****Specification**

5

The present invention relates to recombinant soluble Fc receptors (FcR), recombinant nucleic acids coding for such Fc receptors, host cells containing corresponding nucleic acids as well as a process for the determination of the amount of antibodies of a certain type contained in the blood, plasma or serum of a patient, a process for the determination of the immune status of patients with chronic diseases of the immune system and a process for the screening of substances in view of their ability to act as inhibitors of the recognition and binding of antibodies to the respective cellular receptors. Further, the present invention is concerned with pharmaceutical compositions containing the recombinant soluble FcRs, crystalline preparations of FcRs and FcR/Ig-complexes and especially of the use of such crystalline preparation for the generation of crystal structure data of Fc receptors as well as FcR inhibitors and pharmaceutical compositions containing such FcR inhibitors.

20

A still further subject of the present invention is a recombinant Fc receptor coupled to a solid phase, e.g. a chromatography carrier material. The use of such chromatography material, which is another subject of the present invention, lies in the absorption of immunoglobulins from a body fluid of patients or from culture supernatants of immunoglobulin producing cells.

Fc receptors (FcRs) play a key role in defending the human organism against infections. After pathogens have gained access to the blood circulation they are opsonized by immunoglobulins (Igs). The resulting immunocomplexes bind due to their multivalency with high avidity to FcR bearing cells leading to clustering of the FcRs, which triggers several effector functions (Metzger, H., 1992A). These include, depending on the expressed FcR type and

30

- 2 -

associated proteins, endocytosis with subsequent neutralization of the pathogens and antigen presentation, antibody-dependent cellular cytotoxicity (ADCC), secretion of mediators or the regulation of antibody production (Fridman et al, 1992; van de Winkel and Capel, 1993).

5

Specific FcRs exist for all Ig classes, the ones for IgG being the most abundant with the widest diversity. Together with the high affinity receptor for IgE (FcεRIa), FcγRI (CD64), FcγRII (CD32) and FcγRIIIa (CD16) occur as type I transmembrane proteins or in soluble forms (sFcRs) but also a glycosylphosphatidylinositol anchored form of the FcγRIII (FcγRIIIb) exists. Furthermore, FcγRs occur in various isoforms (FcγRIa, b1, b2, c; FcγRIIa1-2, b1-3, c) and alleles (FcγRIIa1-HR, -LR; FcγRIIIb-NA1, -NA2) (van de Winkel and Capel, 1993). In contrast to the overall homologous extracellular parts, the membrane spanning and the cytoplasmic domains differ. They may be deleted entirely or be of a size of 8 kDa. They may contain either a 26 amino acid immunoreceptor tyrosine-based activation motif (ITAM) as in FcγRIIIa or a respective 13 amino acid inhibitory motif (ITIM) in FcγRIIb involved in signal transduction (Amigorena et al, 1992).

20 Judged by the conserved spacing of cysteins, the extracellular part of the FcRs consists of three (FcγRI, CD64) or two (FcεRI, FcγRII, CD32 and FcγRIII, CD16) Ig-like domains (10 kDa/domain) and therefore belongs to the immunoglobulin super family. These highly glycosylated receptors are homologues, and the overall identity in amino acid sequence among the FcγRs and FcεRIa exceeds 50% in their extracellular regions. Nevertheless, the affinity of FcRs to their ligands varies widely. The higher affinity of  $\approx 10^8 \text{M}^{-1}$  of the FcγRI to Fc-fragment is assigned to its third domain, while the other FcγRs with two domains have an affinity to IgG varying between  $10^5$  and  $10^7 \text{M}^{-1}$ . The affinity of the two domain FcεRIa to IgE exceeds these values by far with a constant of  $10^{10} \text{M}^{-1}$  (Metzger, H., 1992B). In contrast to the mentioned FcRs the low affinity receptor for IgE FcεRII represents a type transmembrane protein and shows a lower homology.

30

- 3 -

FcγRs are expressed in a defined pattern on all immunological active cells. FcγRI is constitutively expressed on monocytes and macrophages and can be induced on neutrophils and eosinophils. The physiological role of FcγRI is still unknown as the expression on monocytes is not vital (Ceuppens et al, 1988). The GPI anchored form of FcγRIII (FcγRIIIb) is exclusively expressed on granulocytes. Due to its missing cytoplasmic part, the signal transduction into the cell occurs solely via other transmembrane proteins like complement receptor type 3 (CR3) that can at least associate with FcγRIIIb (Zhou et al, 1993; Poo et al, 1995). FcγRIIIa is mainly expressed on monocytes and macrophages but only in conjunction with associated proteins (e.g. α- or γ-chains). FcγRII is the receptor with the widest distribution on immunocompetent cells and is mainly involved in the endocytosis of immunocomplexes.

FcγRIIa and FcγRIIb differ in their extracellular region by only 7% of the amino acid residues. Nevertheless, both forms can be distinguished by their binding characteristics to human and mouse IgG subclasses (van de Winkel and Capel, 1993) and their differing affinity to human IgGs (Sondermann et al, 1998A). The situation is rendered even more complicated by the high responder/low responder (HR/LR) polymorphism of FcγRIIa named after the ability of T cells from some individuals to respond to murine IgG1-induced mitogenesis (Tax et al, 1983). Later, it was found that the two exchanges in the amino acid sequence between the LR and the HR form modify the ability to bind human IgG2, which leads to the suggestion that at least one of them is involved in IgG binding (Hogarth et al, 1992).

In contrast to the beneficial role FcRs play in the healthy individual, they also transmit the stimulation of the immune system in allergies (FcεRIa) or autoimmune diseases. Moreover, some viruses employ FcγRs to get access to cells like HIV (Homsy et al, 1989) and Dengue (Littaua et al, 1990) or slow down the immune response by blocking FcγRs as in the case of Ebola (Yang et al, 1998) and Measles (Ravanel et al, 1997).

- 4 -

Hence, the object underlying the present invention was to provide receptors which are easy to produce and can advantageously be used for medical or diagnostic applications. Moreover, it was an object of the invention to provide soluble receptors exhibiting a binding specificity and activity which is analogous to that of the receptors occurring naturally in the human body and which, additionally, make it possible to produce crystals suitable for a structure determination.

This object is accomplished by recombinant soluble Fc receptors which consist only of the extracellular portion of the receptor and are not glycosylated. The receptors according to the present invention are therefore characterized by the absence of transmembrane domains, signal peptides and glycosylation.

Particularly preferred for the present invention are Fc $\gamma$  or Fc $\epsilon$  receptors. This is because IgG and IgE molecules are characteristic for a multiplicity of diseases and conditions, so that their determination and possible ways of influencing them are of great interest. Figure 11 and 12 show an alignment of amino acid sequences of the extracellular parts of some Fc $\gamma$ Rs and Fc $\epsilon$ RI. The FcRs according to the invention include all these sequences or parts thereof that still retain binding capacity to antibodies and/or proper crystallization.

In a particularly preferred embodiment of the invention the recombinant soluble FcR is a Fc $\gamma$ RIIb receptor. Further, it is particularly preferred that the receptor be of human origin. In a particularly preferred embodiment, it contains an amino acid sequence as shown in one of SEQ ID NO:1 to SEQ ID NO:6.

According to the present invention, the preparation of the soluble Fc receptors preferably takes place in prokaryotic cells. After such expression, insoluble inclusion bodies containing the recombinant protein form in

- 5 -

prokaryotic cells, thus facilitating purification by separation of the inclusion bodies from other cell components before renaturation of the proteins contained therein takes place. The renaturation of the FcRs according to the present invention which are contained in the inclusion bodies can principally  
5 take place according to known methods. The advantage of the preparation in prokaryotic cells, the production of inclusion bodies and the thus obtained recombinant soluble Fc receptors make it possible to obtain a very pure and, in particular, also very homogeneous FcR preparation. Also because of the absence of glycosylation the obtained product is of great  
10 homogeneity.

Soluble Fc receptors hitherto produced by recombinant means particularly exhibited the disadvantage that a much more elaborate purification was required, since they were expressed in eukaryotic cells and, due to the  
15 glycosylation which is not always uniform in eukaryotic cells, these products were also less homogeneous.

The recombinant soluble Fc receptors according to the present invention even make it possible to produce crystals suitable for use in X-ray analysis,  
20 as shall be explained later on in the description of further embodiments of the invention. The FcRs of the present invention moreover exhibit practically the same activity and specificity as the receptors naturally occurring in vivo.

A further subject matter of the present invention is a recombinant nucleic  
25 acid having a sequence coding for a recombinant soluble Fc receptor according to the present invention.

The nucleic acid according to the present invention may contain only the coding sequences or, additionally, vector sequences and/or, in particular,  
30 expression control sequences operatively linked to the sequence encoding the recombinant FcR, like promoters, operators and the like.

- 6 -

In a particularly preferred embodiment the nucleic acid of the present invention contains a sequence as shown in one of SEQ ID NO:7 to SEQ ID NO:12. For a comparison, SEQ ID NO:13 and SEQ ID NO:14 show the respective wild type sequences coding for FcγRIIb and FcεRIa. SEQ ID NOs:15-18 show the wild type sequences for FcγRI, FcγRIIa, FcγRIII and FcεRII.

If the nucleic acid of the present invention contains vector sequences, then these are preferably sequences of one or several prokaryotic expression vectors, preferably of pET vectors. Any other known functions or components of expression vectors may also be contained in the recombinant nucleic acid according to the present invention if desired. These may, for instance, be resistance genes allowing for an effective selection of transformed host cells.

A still further subject matter of the present invention is a host cell containing a recombinant nucleic acid according to the present invention. As repeatedly mentioned above, the host cell preferably is a prokaryotic host cell, particularly an E. coli cell.

The recombinant soluble Fc receptors according to the present invention can be used for a multitude of examinations or applications because they specifically react with antibodies. In vivo, the soluble Fc receptors are powerful immunoregulators which, if present in elevated levels, result in a remarkable suppression of the immune system which leads to many partly known and partly not yet understood effects. Based on these effects, several applications of the Fc receptors according to the present invention are further subject matters of the present invention.

One such subject is a process for the determination of the amount of antibodies of a certain type in the blood or serum of a patient, which is characterized by the use of a recombinant soluble FcR according to the



- 7 -

invention in an immunoassay, and the determination of the presence of FcR-antibody complexes. Such assay allows to screen for the presence of a certain kind of antibody and allows also for the determination of the amount of antibodies present in the blood, plasma or serum of a patient.

5

Any type of immunoassay is principally suitable for the use according to the present invention, as long as the presence of FcR-antibody complexes can thereby be detected. Both ELISA (enzyme-linked immunosorbent immunoassay), particularly sandwich assays, and RIA (radio-immunoassay) are suitable, but also competitive testing methods. In a preferred embodiment of the invention where the presence and/or the amount of IgE antibodies is to be examined, an FcεR is used as recombinant soluble receptor according to the present invention. In particular, this method is suited and advantageous for determining a predisposition or manifestation of an allergy.

15

Moreover, a method is preferred in which the presence of soluble FcRs is to be determined and, if required, quantified. For such determination preferably a competitive immunoassay method is used, wherein as competition reagent a recombinant soluble receptor according to the invention is used, most preferably a recombinant FcγR. By means of this test among others the immune status of patients with chronic diseases of the immune system can be determined in a competitive immunoassay. Chronic diseases in the sense of these processes are for instance AIDS, SLE (systemic lupus erythematosus), MM (multiple myeloma) or rheumatoid arthritis, or in the case of FcεRII in B-CLL (Gordon et al., 1987), hyper IgE syndrome (Sarfati et al., 1988) or HCL (Small et al., 1990).

20

25

A further advantageous use of the recombinant receptor according to the present invention lies in the screening of substances in view of their ability to act as inhibitors of the recognition and binding of antibodies to the respective cellular receptors.

30

By means of modern screening techniques such as HTPS (high throughput screening) in combination with multi-well microtiter plates and automatic pipetting apparatuses it is nowadays possible to simultaneously test a multitude of substances for specific properties. As the FcRs according to the present invention can be easily produced at low cost, they can also be  
5 used in such series tests by which substances having an inhibiting effect can easily be identified.

Particularly preferred is such use according to which Fc receptors according to the present invention are used to find or screen inhibitors capable of  
10 inhibiting the recognition and binding of the respective antibodies to the particular receptor of interest.

A further area of application of the substances according to the invention lies in the pharmaceutical field. Hence, a further subject matter of the  
15 invention is a pharmaceutical composition comprising as active agent a recombinant soluble FcR according to the invention. According to the present invention, this pharmaceutical composition may of course comprise conventional useful carrier and auxiliary substances. Such substances are  
20 known to the person of skill in the art, the mode of administration also having to be taken into account. The pharmaceutical composition of the present invention can be advantageously used for the treatment or prevention of autoimmune diseases, allergies or tumor diseases.

25 Soluble forms of Fc receptors such as FcγRIII mediate isotype-specific regulation of B cell growth and immunoglobulin production. In a murine model of myeloma, sFcR suppresses growth and immunoglobulin production of tumor cells (Müller et al, 1985; Roman et al, 1988; Teillaud et al, 1990). Furthermore, sFcR binds to surface IgG on cultures of human IgG-secreting  
30 myeloma cells and effects suppression of tumor cell growth and IgG secretion. Prolonged exposure of these cells to sFcR results in tumor cell cytotoxicity (Hoover et al, 1995).

Also, overreactions of the immune system in allergic reactions or due to massive antigen load might be reduced by, for example, intravenous application of soluble FcR (Ierino et al, 1993).

5 Therefore, a preferred pharmaceutical composition according to the invention for use in the treatment of AIDS, rheumatoid arthritis or multiple myeloma contains a recombinant soluble Fc $\gamma$  receptor and, preferably, a receptor having the amino acid sequence as shown in SEQ ID NO:1-4.

10 It was also of great interest to obtain crystal structure data of Fc receptors and/or Fc receptor/Ig complexes. On the one hand, these are a key to the understanding of molecular mechanisms in immunocomplex recognition. On the other hand, these structural data can be used to find out common features in the structures of different Fc receptors and use the knowledge  
15 of the structures to generate inhibitors or identify and produce new artificial antibody receptors.

It was also of great interest to obtain information on the concrete binding sites of immunoglobulins to their respective receptors in naturally occurring  
20 three-dimensional molecules. Therefrom even more precise findings on the interactions between antibody and receptor can be obtained and also on how these interactions can be modulated. In this connection modulation means either an enhancement of the interaction or a reduction leading to an inhibition by e.g. covering the binding sites on one or more parts of the  
25 complex.

To obtain such crystal structure data and conformation information, a crystalline preparation of the recombinant soluble Fc receptor according to the invention is used. The recombinant soluble FcRs according to the  
30 invention surprisingly can be obtained pure enough to produce crystals that give reliable X-ray structure determination data. Such crystallization was not

- 10 -

possible with the hitherto produced receptor molecules, mostly due to their lack of homogeneity.

Therefore, another embodiment of the present invention concerns a  
5    crystalline preparation of an Fc receptor according to the invention. Yet  
another embodiment of the present invention is a crystalline preparation of  
a complex of soluble Fc receptor according to the invention together with  
the related immunoglobulin Fc part. Particular preferred embodiments are  
shown in the examples as well as the relevant crystal structure data. Via  
10   crystal structure analysis of the crystalline preparations the exact amino  
acids of the Fc receptor/Ig complexes could be detected which mediate the  
coupling. These amino acids are shown in Fig 6a and 6b and the type of  
binding between the individual amino acids of both molecules in the  
complex is also indicated. A further embodiment of the present invention is  
15   therefore the use of a crystalline preparation of a recombinant soluble Fc  
receptor for the generation of crystal structure data of Fc receptors. From  
this crystal structure data information about the three-dimensional structure  
and the active sites for the binding of antibodies can be obtained. Especially  
preferably is the use of a crystalline preparation of a complex of  
20   recombinant soluble Fc receptor according to the invention and the  
corresponding immunoglobulin molecule for the generation of crystal  
structure data for the complexes. These data allow to determine the actual  
interactions that are formed between the two molecules and allow for the  
first time to obtain exact information about the interaction of the molecules  
25   thereby conferring knowledge about possible sites for inhibition or  
enhancement of the binding. On the basis of the information obtained from  
the crystal structure data the findings necessary for effecting modulation of  
the interaction between Fc receptor and immunoglobulin can be obtained.  
This modulation can range from enhancement to complete inhibition to  
30   an inhibition of the binding.

- 11 -

The stated applications are merely preferred embodiments of the use of the crystal structure data. Many other applications seem possible, too.

5 Suitably, the structural data for the generation and/or identification of inhibitors or new receptors, respectively, are used in a computer-aided modelling program.

Particularly preferred for the present invention are the structures of FcRs or FcR:Fc-fragment complexes as exemplified in figures and examples. Such  
10 structures can be used to design inhibitors, antagonists and artificial receptor molecules.

Computer programs suitable for computer-aided drug design and screening are known to the person skilled in the art and generally available. They  
15 provide the possibility to examine umpteen compositions on the computer in view of their ability to bind to a certain molecule when the corresponding structure dates are entered in the computer. With the help of this possibility a great number of known chemical compositions can be examined regarding their inhibiting or antagonistic effect. The person skilled in the art merely  
20 requires the crystal structure dates provided by the present invention and a commercially available screening program (Program Flexx: From the GMD-German National Research Center for Information Technology, Schloss Birlinghoven, D-53754 Sankt Augustin, Germany). A preferred embodiment of the present invention therefore is the use of the crystal structure data  
25 obtained for the recombinant soluble Fc receptor according to the invention and for the complexes of recombinant soluble Fc receptor according to the invention and corresponding immunoglobulin in a computer aided modelling program for the identification and production of Fc receptor inhibitors.

30 Likewise, a further embodiment of the present invention is the use of the crystal structure data obtained for the receptors according to the invention and the receptor/immunoglobulin complexes, respectively for the

- 12 -

identification and preparation of new Fc receptors which can be used, e.g. as antagonists and competitors. The crystal structure data and the data on the amino acids involved in the binding to Fc receptors obtained therefrom can serve for example to generate mutated immunoglobulins which can also be used as inhibitors. It is imaginable that mutated or chemically modified inhibitors undergo tight binding and thus effect a blocking of receptors. On the other hand, the data obtained for the binding sites of immunoglobulins can also be used for the identification and/or preparation of inhibitors for immunoglobulin molecules. Since the present invention teaches the binding sites to the receptor, it is easy to effect a blocking of the binding sites with the help of relatively simple molecules. Therefore, a further subject matter of the present invention is the use of the crystal structure data obtained for the FcR/Ig complexes for the identification and/or preparation of immunoglobulin inhibitors.

15

Accordingly, still further subject matter of the present invention are FcR inhibitors which have a three-dimensional structure which is complementary to the recombinant soluble FcR according to the invention and inhibit the binding of antibodies to FcRs.

20

Another further subject of the present invention are immunoglobulin inhibitors which have a three-dimensional structure which is complementary to the immunoglobulin binding site for recombinant soluble Fc receptors according to the invention and inhibit the binding of immunoglobulins to Fc receptors.

25

The term "complementary" is to be understood within the framework of the invention in such a way that the inhibitor molecules must be substances which are able to cover at least so many binding sites on the immunoglobulin or on the Fc receptor that the binding between Fc receptor and immunoglobulin is at least decisively weakened. Covering can take place both by binding to the amino acids mediating the complex formation

30

- 13 -

of either component but also in such a way that at least complex formation is no longer possible, be it by sterically inhibition or by binding to adjacent amino acids, however, covering the amino acid involved in the complex binding between Fc receptor and immunoglobulin.

5

In connection with the present invention it was possible for the first time to determine the exact binding sites and the amino acids involved in the binding of the antibody and antibody receptor molecules. One is now able to design specifically binding molecules and to screen candidate compositions on the computer. This enables the selection of such compositions from a variety of possibly candidate compositions which can effect a sufficient inhibition of complex formation between Fc receptor and immunoglobulin.

15 What is important for the inhibitors of the invention is that, owing to their structure and specificity, they are capable of binding to the FcRs or immunoglobulins and thus prevent the normal binding between FcRs and the constant parts of antibodies.

20 Preferably, such FcR or IgG inhibitors are small organic molecules which can easily be administered orally. They are an interesting alternative to cortisone in the treatment of autoimmune diseases and host/graft rejections. Such a molecule would also suppress reinfection rates with certain viruses, e.g. Dengue virus where the antibody coated virus is Fc $\gamma$ RIIb dependent internalized (Littaua et al, 1990), HIV where on CD4 positive T cells an antibody enhancement of HIV infection is mediated by Fc $\gamma$ RIII (Homsy et al, 1989), or Ebola where the virus secreted glycoprotein inhibits early neutrophil activation by blocking sFc $\gamma$ RIII which affects the host response to infection (Yang et al, 1998).

30

The development of inhibitors also leads to substances that interfere with the recognition of IgE by their receptors. From the modelled structure of

- 14 -

FcεRI, peptides have already been developed which inhibit mast cell degranulation in vitro. With the now available knowledge of the structures of the homologue receptors and the receptor-antibody complex in atomic detail, a new possibility for a rational drug design is opened.

5

The Fc-receptor bind between the two CH2-domains of the Fc-fragment in the so-called lower hinge region (Fig.8). The binding region of the Fc-receptor is described in Example 1 (The contact interface to IgG). The residues promoting the interaction between FcR and immunoglobulin are shown in figures 7, 10a and 10b. Thereby three interaction regions become evident (Fig.5).

10

**1st region: FcR (residues 85 to 87 and residue 110) - Ig (Chain A residues 326-328)**

15

Proline 328 of the Ig is clamped by the residues Trp 87 and 110 in a sandwich like manner. These residues are conserved among the IgG and IgE receptors as well as in the IgG and IgE. An inhibitor binding to this prominent region would strongly interfere with binding. This region is additionally attractive for inhibitor design because the exposed hydrophobic surface region comprising the residues Trp 87, Ile 85, Gly 86 of the receptors could be employed to obtain additional binding energy. The functional groups of Thr 113 and Glu 18 and Lys 19 side chains in the vicinity may contribute especially to specific inhibitor binding.

20

**2nd region: FcR (residues 126-132 and residues 155-158) - Ig (Chain A and Chain B residues 234-239)**

25

The amino terminal residues 234-239 of both Ig chains are recognised differently by the FcR, thereby breaking the 2-fold symmetry of the Fc fragment.

30

This residues of Fc-fragment chain A are in contact with residues Val 155 - Lys 158 of the receptor and the same residues from Fc-fragment chain B with receptor residues Gly 126 - His 132. This region shows the most



- 15 -

differences in the sequence alignment of the receptors as well as the immunoglobulins and should therefore be involved in specificity generation. This deep cleft between the Fc-fragment chains is well suited for inhibitor design and would be the site of choice for the development of inhibitors when issues of specificity are concerned.

**3rd region: FcR (residues 117, 126 and 129-132) - Ig (Chain B residues 264-265 and residues 296-297)**

This binding region is characterised by a clustering of amino acid residues carrying functional groups in their side chains, that might be employed in various ways for inhibitor design on the receptor and the Ig side of the contact.

Molecules that interact with one or more of the above described regions, and are designed or screened explicitly for exploiting the knowledge of binding sites are considered as inhibitors according to the invention.

Further subject matters of the present invention are pharmaceutical compositions containing as active agent an FcR inhibitor or an immunoglobulin inhibitor as mentioned above. Such pharmaceutical compositions may, for example, be used in the treatment or prevention of diseases which are due to overreactions or faulty reactions of the immune system, preferably the treatment or prevention of allergies, autoimmune diseases or anaphylactic shock.

A further subject of the present invention is the sFcR according to the invention, bound to a solid phase. Such heterogeneous receptors can be used for immunoassays or other applications where the receptor in an immobilized form can be used beneficially.

In a preferred embodiment of the invention the solid phase is a chromatography carrier material onto which the Fc receptor is fixed, e.g.

- 16 -

sepharose, dextran sulfate etc. Such chromatography materials with Fc receptors bound thereto can beneficially be used for the adsorption of immunoglobulins from the blood, plasma or serum of patients or from the culture supernatant of immunoglobulin producing cells (meaning  
5 concentration, enrichment and purification of antibodies).

On the one hand, the antibodies bound to the chromatography material can be eluted and, for example, the immune status of a patient can thereby be determined. On the other hand, antibodies from the blood of a patient can  
10 thereby be enriched before carrying out further tests, which is a further preferred embodiment of the present invention. In many cases it is difficult to conduct diagnostic assays using blood samples if the latter contains only a very small number of the antibodies to be identified. By means of a concentration using a specific chromatographic column with Fc receptors  
15 according to the present invention, antibodies of interest can easily be concentrated and separated from many other substances which might disturb the test.

Basically, it is also possible to use a chromatography material according to  
20 the present invention in an extracorporeal perfusion system for lavage of the blood in case of certain diseases where the removal of antibodies plays a crucial role.

It is, however, also possible to use another material as solid phase to which  
25 the soluble Fc receptor according to the invention is coupled, e.g. microtiter plates or small reaction vessels to the walls of which Fc receptors are bound either directly or indirectly. Such solid phases and vessels can be particularly important for diagnostic methods, as they enable screening by using immunoassays e.g. for detecting the presence of certain  
30 immunoglobins in patients' blood or other body fluids.

To sum up, the recombinant soluble Fc receptors provided by the present invention as well as the corresponding structure determination of crystalline

- 17 -

preparations of these receptors and of crystalline complexes of receptors and immunoglobins enable for the first time to perform a rational drug design, wherefrom it is possible to modulate the interaction between immunoglobulins and Fc receptors on cells or soluble receptors. Such a modulation is preferably an inhibition, whereby the inhibition of the formation of a complex from IgG and Fc receptor takes place by covering and preferably by binding of inhibitor molecules to the Fc receptor or the immunoglobulin. There are various medical applications for such modulating drugs and in particular of inhibitors and only few of these applications have been exemplary mentioned within the framework of the present specification. This can and should by no means exclude the applicability of such molecules which have been designed or screened on the basis of the findings about the molecular structure or FcR/Ig complexes disclosed herein for the treatment or prevention of other health disturbances.

The following Examples are to further illustrate the invention in conjunction with the Figures.

### **Example 1**

#### **shFcγRIIb (soluble human FcγRIIb)**

##### **1.1 Cloning and Expression**

The cDNA of human FcγRIIb2 (Engelhardt et al, 1990) was modified using mutagenous PCR (Dulau et al, 1989). Therefore, a forward primer was used for the introduction of a new start methionine after the cleavage site of the signal peptide within a *Nco*I site (5'-AAT AGA ATT CCA TGG GGA CAC CTG CAG CTC CC-3') while the reverse primer introduced a stop codon between the putative extracellular part and the transmembrane region followed by a *Sa*I site (5' CCC AGT GTC GAC AGC CTA AAT GAT CCC C-3'). The PCR product was digested with *Nco*I and *Sa*I, cloned into a pET11d expression vector (Novagen) and the proposed sequence was

- 18 -

confirmed. The final construct was propagated in BL21(DE3) (Grodberg and Dunn, 1988). For the overexpression of FcγRIIb a single colony of the transformed bacteria was inoculated in 5ml LB medium containing 100 μg ampicillin per ml (LB-Amp100) and incubated overnight at 37°C. The culture was diluted 200-fold in LB-Amp100 and incubation was continued until an OD600 of 0.7-0.9 was achieved. The overproduction of the protein was induced by adding IPTG to a final concentration of 1 mM. After a growing period of 4 hours the cells were harvested by centrifugation (30 min, 4000 x g) and resuspended in sonification buffer (30 mM sodium phosphate, 300 mM sodium chloride, 0.02% sodium azide, pH 7.8). After addition of 0.1 mg lysozyme per ml suspension and incubation for 30 min at room temperature the sonification was performed on ice (Branson Sonifier, Danbury, CT; Macrotip, 90% output, 80% interval, 15 min). The suspension was centrifuged (30 min, 30,000 x g) and resuspended with a Dounce homogenizer in sonification buffer containing 0.5% LDAO. The centrifugation step and resuspension in LDAO containing buffer was repeated once before this procedure was repeated twice without LDAO. The purified inclusion bodies were stored at 4°C.

## 1.2 Refolding and purification of soluble human FcγRIIb (shFcγRIIb)

The purified inclusion bodies were dissolved to a protein concentration of 10 mg/ml in 6 M guanidine chloride, 100 mM 2-mercaptoethanol and separated from the insoluble matter by centrifugation. The refolding was achieved by rapid dilution. Therefore, one ml of the inclusion body solution was dropped under stirring within 15 hours into 400 ml of the refolding buffer (0.1 M TRIS/HCl, 1.4 M arginine, 150 mM sodium chloride, 5 mM GSH, 0.5 mM GSSG, 0.1 mM PMSF, 0.02% sodium azide, pH 8.5, 4°C). Afterwards, the mixture was stirred for 2-3 days until the concentration of free thiol groups was reduced to 1 mM by air oxidation as measured according to Ellman (Ellman, 1959). The solution was dialyzed against PBS and sterile filtered before it was concentrated 10-fold in a stirring cell

- 19 -

equipped with a 3kD MWCO ultrafiltration membrane. The protein solution was applied to a hlgG sepharose column (50 mg hlgG per ml sepharose 4B). Unbound protein was washed out with 50 mM TRIS pH 8.0 before elution of FcγRIIb by pH jump (150 mM sodium chloride, 100 mM glycine, 0.02% sodium azide, pH 3.0). The eluate was immediately neutralized with 1 M TRIS pH 8.0. The FcγRIIb containing solution was concentrated and subjected to gel filtration on a Superdex-75 column equilibrated with crystallization buffer (2 mM MOPS 150 mM sodium chloride, 0.02% sodium azide pH 7.0). The fractions containing FcγRIIb were pooled, concentrated to 7 mg/ml and stored at -20°C.

### 1.3 Equilibrium gel filtration experiments

A Superdex75 column was connected to FPLC and equilibrated with PBS containing 10 μg shFcRIIb per ml. Human Fc fragment was solved to a concentration of 1 μg/10 μl in the equilibration buffer and injected. The resulting chromatogram yielded a positive peak comprising the complex of the shFcγRIIb and the Fc fragment while the negative peak represents the lack of receptor consumed from the running buffer for complex formation.

### 1.4 Crystallization and data collection

Initial crystallization trials employing a 96 condition sparse matrix screen (Jancarik and Kim, 1991) were performed in sitting drops at 20 °C using the vapor diffusion method. Occuring crystals were improved by changing the pH as well as the salt, precipitant and additive concentration. Diffraction data from suitable crystals was collected on an image plate system (MAR research) using graphite monochromated CuK<sub>α</sub> radiation from a RU200b rotating anode generator (Rigaku) operated at 50 kV and 100 mA. The reflections were integrated with the program MOSFLM (Leslie, 1997) and subsequently the data was scaled, reduced and truncated to obtain the

- 20 -

structure-factor amplitudes using routines from the CCP4 program suite (Collaborative Computational Project, 1994).

### 1.5 Summary of expression, purification and refolding of shFcγRIIb

5

The extracellular part of FcγRIIb was expressed in high levels under the control of a T7 promoter in the T7 RNA polymerase positive *E. coli* strain BL21/DE3 (Grodberg & Dunn, 1988). The protein was deposited in inclusion bodies, which were employed in the first purification step. The isolation of the inclusion bodies was started with an intense combined lysozyme/sonification procedure to open virtually all cells which would otherwise contaminate the product. The subsequent washing steps with the detergent LDAO, which has excellent properties in solubilizing impurities but not the inclusion bodies itself already yielded a product with a purity of > 90% (Fig. 1).

15

This product was used for refolding trials without further purification. The inclusion bodies were dissolved in high concentration of 2-mercaptoethanol and guanidine to ensure the shift of covalent and non-covalent aggregates to monomers. This solution was rapidly diluted with refolding buffer to minimize contacts between the unfolded protein molecules which would otherwise form aggregates. The use of arginine in the refolding buffer prevents the irreversible modification of side chains as often recognized with urea. After addition of the protein to the refolding buffer, the solution was stirred at 4 °C until the concentration of free thiol groups was reduced to 1 mM, which was absolutely necessary as earlier dialysis resulted in an inactive product. In a second purification step the dialyzed and refolded FcγRIIb was bound to immobilized hIgG to remove minor fractions of *E. coli* proteins and inactive receptor. The protein was eluted with a pH jump and immediately neutralized. After this affinity chromatography step shFcγRIIb is essentially pure except for a minor contamination resulting from the coeluting IgG which leached out of the matrix even after repeated use (Fig.

20  
25  
30

- 21 -

- 1). The IgG as well as receptor multimers which are not visible in the reducing SDS-PAGE could easily be removed by gel filtration. Parallel to the removal of the contaminants in this step the buffer is quantitatively exchanged. This procedure ensures a defined composition of the protein solution as even slight variations can cause irreproducibility of the crystallization attempts or even inhibit the formation of crystals. Overall 6 mg pure protein could be gained per litre E. coli culture, which is about 10 % from the FcγRIIb content of the inclusion bodies.
- 10 N-terminal protein sequencing revealed the identity with the expected sequence H<sub>2</sub>N-GTPAAP without detectable contamination. ESI-MS analysis showed that the final material used in crystallization trials is homogenous with respect to size. From the primary sequence the molecular weight was calculated to 20434 Da, which corresponds to 20429 Da found by mass spectroscopy. The discrepancy lies within the error of the instrument, and no additional peak for a species containing the leading methionine is found.
- 15

The crystallization of shFcγRIIb was performed in sitting drops using the vapor diffusion method. Initial trials with a sparse matrix screen (Jancarik & Kim, 1991) resulted already in small crystalline needles. Subsequent optimization of the preliminary crystallization condition by varying precipitant, salt, their concentration and pH led to the isolation of three different crystal forms. Orthorhombic crystals grew from mixture of 1.5 μl reservoir solution (33% PEG2000, 0.2 M sodium acetate, pH 5.4) with 3 μl of the protein solution. They appeared within 3 days and reached their final size of approximately 80 μm x 80 μm x 500 μm after one week. These crystals diffracted to 1.7 Å. Crystals could also be grown in two other space groups from reservoir solution containing 26% PEG8000, 0.2 M sodium acetate, pH 5.6, 5 mM Zn(OAc)<sub>2</sub>, 100 mM sodium chloride (hexagonal form) and 26% PEG8000, 0.2 M NaOAc, pH 5.6, 10% (v/v) 1,4-Dioxan, 100 mM sodium chloride (tetragonal form). These crystals were

20

25

30

- 22 -

of suitable size for X-ray analysis but diffracted only to 2.7 Å and 3.8 Å for the tetragonal and hexagonal crystal form respectively (Table 1).

FcγRII was expressed in *E. coli* which, besides the comparatively low  
5 production costs and the availability, has several advantages especially  
when the glycosylation performed by mammalian cells is not necessary for  
the function of the protein as in the case of FcγRII where IgG binding  
occurs independently of carbohydrate attachment (Sondermann et al,  
1998A). In *E. coli* a homogenous product can reproducibly be generated,  
10 which is in contrast to the expression in mammalian cells where batch  
dependent variances are often observed. In such a system the product is for  
several days exposed to proteases at temperatures of more than 30°C. In  
contrary, the expression of the protein in *E. coli* under the control of the  
strong T7 promoter at 37°C frequently leads to the formation of protease  
15 inaccessible inclusion bodies. A further advantage of the expression in  
bacteria is that the material could be considered to be free of pathogenic  
germs, which might derive from employed fetal calf serum or the cell line  
itself. In mammalian expression particular care must be taken during the  
purification of the target protein because potential effective hormones or  
20 growth factors might be copurified. One case where the effects of sFcγR  
were ascribed to a TGFβ1 contamination is already reported (Galon et al,  
1995).

## 1.6 Purification

25

The purification procedure is straightforward. It consists of three steps  
which can easily be performed in a single day. The protein is obtained in a  
pure form and in high yields and could even be obtained in considerable  
quality without the expensive IgG affinity column. The success of such a  
30 protocol would depend on the careful preparation of the inclusion bodies,  
as most of the impurities can be eliminated already in the first purification  
step.



## 1.7 Characterization

The purified FcγRIIb was characterized by SDS-PAGE and isoelectric focussing as well as N-terminal sequencing and mass spectroscopy. Thus, the material can be considered pure and homogeneous with respect to its chemical composition, but the intriguing question whether the receptor is correctly folded remains to be discussed. All cysteins are paired, since no free thiol groups are detected with Ellman's test. The material is monomeric and eludes with the expected retention time in peaks of symmetrical shape from a size exclusion chromatography column. Furthermore, FcγRIIb binds to IgG sepharose, recombinant FcγRIIb from E. coli is active because it specifically binds IgG.

## 1.8 Crystallization

The orthorhombic crystal form of FcγRIIb diffracted X-rays to a resolution of 1.7 Å, which is a drastic improvement compared to previously reported crystals of the same molecule derived from insect cell expression (Sondermann et al, 1998A). These crystals diffracted to 2.9 Å and were of space group P3<sub>1</sub>21. Thus, the glycosylation of the insect cell derived receptor influences the crystallization conditions. Instead of the trigonal space group, three different crystal forms are found. After a possible solution of the structure these crystal forms will help identify artificial conformations of the protein due to crystal contacts.

FcγRs do not exhibit any sequence similarity to other proteins but due to a conserved cystein spacing they are affiliated to the immunoglobulin super family. Consequently, we tried to solve its structure by molecular replacement, but extensive trials using IgG domains from a variety of molecules failed. Thus the structure of FcγRIIb has to be solved by the methods of multiple isomorphous replacement.

- 24 -

We have shown for the first time that FcγRIIb can be obtained in an active form from *E. coli*. This is the basis for crystallographic investigations that will soon, due to the already gained crystals of exceptional quality, result in the structure solution of this important molecule. The structure will provide information on the IgG binding site and provide a starting point for the knowledge based design of drugs that interfere with recognition of the ligand by its receptor. Furthermore, because of the high homology between FcγRIIb and other FcRs including FcεRIa it seems possible that these molecules can be produced in the same way, which would provide valuable material for the ongoing research.

## 1.9 Methods

### Protein chemistry

Recombinant soluble human FcγRIIb was expressed in *E. coli*, refolded purified and crystallized as described elsewhere (Sondermann et al, 1998B). Briefly, the putative extracellular region of hFcγRIIb2 (Engelhardt et al, 1990) was overexpressed in *E. coli*. Inclusion bodies were purified by lysozyme treatment of the cells and subsequent sonification. The resulting suspension was centrifuged (30 min 30,000 x g) and washed with buffer containing 0.5% LDAO. A centrifugation step and resuspension in LDAO containing buffer was repeated once before this procedure was repeated twice without LDAO. The inclusion bodies were solved in 6 M guanidine hydrochloride and the protein was renatured as described. The dialyzed and filtrated protein solution was applied to a hIgG sepharose column and eluted by pH jump. The concentrated neutralized fractions were subjected to size-exclusion chromatography on a Superdex-75 column (26/60, Pharmacia).

### Crystallization

Crystallization was performed in sitting drops at 20°C using the vapor diffusion technique. Crystallization screens were performed by changing pH,

- 25 -

salt, precipitant and additives. The final crystals used for data collection were grown in 33% PEG2000, 0.2 M sodium acetate, pH 5.4 (orthorhombic form) 26% PEG8000, 0.2 M sodium acetate, pH 5.6, 10% (v/v) 1,4-dioxane, 100 mM sodium chloride (tetragonal form), and 26%  
5 PEG8000, 0.2 M sodium acetate, pH 5.6, 5mM ZN(OAc)<sub>2</sub>, 100 mM sodium chloride (hexagonal form). The insect cell derived protein was crystallized in 32% PEG6000, 0.2 M sodium acetate, pH 5.3.

#### Preparation of heavy-atom derivatives

10 The heavy-atom derivatives were prepared by soaking the crystals in the crystallization buffer containing 2 mM platinum(II)-(2,2'-6,2''terpyridinium) chloride for 24 hours or 10 mM uranylchloride for 8 days.

#### X-ray data collection

15 Diffraction data was collected on an image plate system (MAR research) using graphite monochromated CuK<sub>α</sub> radiation from a RU200b rotating anode generator (Rigaku) operated at 50 kV and 100 mA. The reflections were integrated with the program MOSFLM 5.50 (Leslie, 1997) and subsequently the data was scaled and truncated to obtain the  
20 structure-factor amplitudes using routines from the CCP4 program suite (Collaborative Computational Project, 1994).

#### Structure determination

The structure was solved with the standard procedures of the MIR method.  
25 From the large number of soaks carried out with different heavy-atom components only the two compounds yielded interpretable Patterson maps. The heavy-atom positions for each derivative were determined from difference Patterson maps and initial phases were calculated. Cross-phased difference Fourier maps were used to confirm heavy atom positions and  
30 establish a common origin for the derivatives. Anomalous data were included to discriminate between the enantiomers. The heavy atom parameters were further refined with the program MLPHARE from the CCP4

- 26 -

package leading to the statistics compiled in Table 2. An electron-density map was calculated to a resolution of 2.1 Å and the phases were improved further by solvent flattening and histogram matching with the program DM from the CCP4 suite. The resulting electron density map was of sufficient quality to build most of the amino acid residues. Model building was performed with O (Jones et al, 1991) on an Indigo2 work station (Silicon Graphics Incorporation). The structure refinement was done with XPLOR (Brünger et al, 1987) by gradually increasing the resolution to 1.7 Å using the parameter set of Engh and Huber (Engh & Huber, 1991). When the structure was complete after several rounds of model building and individual restraint B-factors refinement ( $R_{\text{fac}} = 29\%$  /  $R_{\text{Free}} = 36\%$ ), 150 water molecules were built into the electron density when a Fo-Fc map contoured at 3.5  $\sigma$  coincided with well defined electron density of a 2Fo-Fc map contoured at 1  $\sigma$ . The resulting refinement statistic is shown in Table 3.

### 1.10 Structure determination

The crystal structure of recombinant soluble human FcγRIIb was solved by multiple isomorphous replacement (MIR) to 1.7 Å resolution, since a structure solution by molecular replacement with isolated domains of the Fc fragment from human IgG1 (Huber et al, 1976, PDB entry 1fc1; Deisenhofer, 1981) failed. The putative extracellular part of the receptor (amino acid residues 1-187 as depicted in SEQ ID NO:2) was used for crystallization trials (Sondermann et al, 1998B) while the model contains the residues 5-176 as the termini are flexible and not traceable into the electron density. Additionally, the model contains 150 water molecules and the refinement statistics are summarized in Table 2. The structure contains a cis proline at position 11. None of the main chain torsion angles is located in disallowed regions of the Ramachandran plot. The fully refined model was used to solve the structure of the same protein in crystals of space group P4<sub>2</sub>2<sub>1</sub>2 and of the glycosylated form derived from insect cells in crystals of space group P3<sub>1</sub>21 (Table 2).

- 27 -

The polypeptide chain of FcγRIIb folds into two Ig-like domains as expected from its affiliation with the immunoglobulin super family. Each domain consists of two beta sheets that are arranged in a sandwich with the conserved disulfide bridge connecting strands B and F on the opposing sheets (Fig. 3). Three anti-parallel β-strands (A1, B, E) oppose a sheet of 5 β-strands (C', C, F, G, A2), whereby strand A1 leaves the 3-stranded β-sheet and crosses over to the 4-stranded anti-parallel sheet adding the short parallel 5th strand A2. The arrangement of secondary structure elements as well as their connectivity is identical in both domains of the FcγRIIb and a rigid body fit of one domain onto the other revealed a r.m.s. distance of 1.29 Å of 67 matching Cα atoms.

The domains are arranged nearly perpendicularly to each other enclosing an angle of 70 degrees between their long axes forming a heart-shaped overall structure. This arrangement results in an extensive contact region between the domains (Fig. 4). Residues from strand A2 and from the segment linking A2 and A1 of the N-terminal domain intermesh with residues of strands A1 and B from the C-terminal domain. This region is tightly packed and the interaction is strengthened by several hydrogen bonds resulting in a rigid arrangement. This is confirmed by the conservation of the structure in three different space groups. In orthorhombic, tetragonal and hexagonal (insect cell derived) crystal forms a deviation of less than 2° in the interdomain angle is found.

### 1.11 Overall structures

The structure of recombinant human FcγRIIb derived from E.coli was solved by MIR to 1.7 Å resolution from orthorhombic crystals. An essentially identical structure is found in tetragonal and with protein derived from insect cells in hexagonal crystals. In all three structures the last nine residues of the polypeptide chain were found disordered. The flexibility of the C-terminal linker region between the structured core of the molecule and

- 28 -

the transmembrane part may be functionally relevant to allow some reorientation of the receptor to enhance the recognition of the Fc parts in immunocomplexes.

## 5 1.12 Homologue receptors

The Ig domains found in the Ig super family of proteins are characterized by a beta sandwich structure with a conserved disulfide bridge connecting two strands of the opposing sheets. The typical arrangement of 3 and 4 anti  
10 parallel beta strands that form a sandwich as found in FcγRIIb occurs also in the T cell receptor, Fc fragment, CD4 or the Fab fragment. A structural alignment of the individual Ig domains of these molecules with the two domains of FcγRIIb shows a common, closely related structure. The relative arrangement of the domains, however, is not related in these molecules and  
15 covers a broad sector. Despite the structural similarity between Ig domains from different molecules and the strikingly low r.m.s. deviation of Cα atoms that result when the two domains of FcγRII are superimposed, no significant sequence similarity is found (Figs. 5a and 5b). A structure-based sequence alignment shows a conserved hydrophobicity pattern along the sequence  
20 of the domains, together with, beside the cysteins, only few identical amino acid residues. We first prepared a structure-based alignment of the two C-terminal domains of the IgG1 heavy chain and the FcγRIIb and added the sequences of the other related FcγRI and the FcεRIa domains. This shows that the sequences of the three domain FcγRI and the two domain receptors  
25 are compatible with the hydrophobicity pattern of Ig domains and several conserved amino acid residues are revealed. Firstly, the different domains of an FcR are more related to each other than to Ig domains from other molecules of the Ig super family. Secondly, the N-terminal domains of the receptors relate to each other as the second domains do. Thirdly, the  
30 sequence of the third domain of FcγRI shows features from both groups of domains. Taken together, we confirm the affiliation of the FcRs to the Ig super family and speculate that all FcR-domains originate from a common

ancestor, an ancient one domain receptor that acquired a second domain by gene duplication. Further divergent development of such a two domain receptor resulted in the present diversity, including FcγRI that acquired a third domain.

5

Conservation of these amino acid residues that contribute to the interdomain contact in FcγRIIb in the alignment are a hint to a similar domain arrangement in different receptors. In Table 4 the residues contributing with their side chains to the interdomain contact (Fig. 4) are compiled for FcγRIIb together with the corresponding amino acid residues in other receptors according to the structure-based sequence alignment of Fig. 5b. Except for Asn15, which is not conserved between the FcRs, the involved residues are identical or conservatively replaced providing strong support for a similar structure and domain arrangement in all FcRs.

15

### 1.13 The contact interface to IgG

Limited information about the interactions of FcRs with their ligands is available from mutagenesis studies (Hogarth et al, 1992; Hulett et al, 1994; Hulett et al, 1995). By systematically exchanging loops between the β-strands of FcγRIIa for FcεRIa amino acid residues the B/C, C'/E and F/G loops of the C-terminal domain were evaluated as important for ligand binding (Fig. 3, Fig. 5b). In the structure model these loops are adjacent and freely accessible to the potential ligand. Additionally, most of the amino acid residues in these loops were exchanged for alanines by single site mutations which resulted in a drastic alteration of the affinity of FcγRIIa to dimeric human IgG1. Also, the single amino acid exchange Arg 131 to His in the C-terminal domain (C'/E loop) in the high responder/low responder polymorphism, which alters the affinity of the FcγRIIa to murine IgG1, points to that region. Thus, the amino acid residues in this area are either important for ligand binding or the structural integrity of that region. Here, the structure shows a clustering of the hydrophobic amino acid residues Pro

30

- 30 -

114, Leu 115 and Val 116 in the neighbourhood of Tyr 157. This patch is separated from the region Leu 159, Phe 121 and Phe 129 by the positively charged amino acid residues Arg 131 and Lys 117 which protrude from the core structure (Fig. 5b).

5

### 1.14 Glycosylation

In the sequence of FcγRIIb three potential N-glycosylation sites are found. All three sites are on the surface of the molecule and are accessible. They are located in the E/F loops (N61 and N142) of both domains and on strand E (N135) of the C-terminal domain (Fig. 3, Fig. 6). Since the material used for the solution of this structure was obtained from *E. coli*, it does not contain carbohydrates, while the FcRs isolated from mammalian cells are highly glycosylated. The three potential glycosylation sites are located rather far from the putative IgG binding region, and non-glycosylated FcγRIIb binds human IgG, suggesting a minor role of glycosylation in binding. This was confirmed by the structure of the FcγRIIb produced in insect cells which is glycosylated (Sondermann et al, 1998A). Except for a 2° change of the interdomain angle possibly due to different crystal contacts, no differences between the glycosylated and unglycosylated protein structures were found. The three glycosylation sites are only optionally used as shown by SDS-PAGE where the material appears in 4 bands. No additional electron density for those sugars was found a consequence of chemical and structural heterogeneity.

25

### Example 2

#### **shFcγRIIa (soluble human FcγRIIa)**

The procedures were performed according to example 1 except for the indicated changes:

30



## 2.1 Cloning and Expression

shFcγRIIIa was generated by mutating the respective wild-type cDNA (Stengelin et al., 1988) and expressed according to example 1 with the mutagenous primers listed in table 5. For the expression of the protein a pET22b + vector was chosen.

## 2.2 Refolding and purification

shFcγRIIIa was refolded according to example 1 with the respective refolding buffer listed in table 6.

## 2.3 Crystallisation

shFcγRIIIa was crystallised as described under conditions indicated in table 7.

## 2.4 Structure determination

The structure was solved with the method of isomorphous replacement with shFcγRIIb as search model.

## Example 3

### shFcγRIII (soluble human FcγRIII)

The procedure was performed according to example 1 except for the indicated changes:

## 3.1 Cloning and Expression

shFcγRIII was generated by mutating the respective wild-type cDNA (Simmons & Seed, 1988) and expressed according to example 1 with the mutagenous primers listed in table 5. For the expression of the protein a pET22b + vector was chosen.

### 3.2 Refolding and purification

shFcγRIII was refolded according to example 1 with the respective refolding buffer listed in table 6.

### 5 3.3 Crystallisation

shFcγRIII was crystallised as described under conditions indicated in table 7.

### 3.4 Structure determination

10 The structure was solved with the method of isomorphous replacement with shFcγRIIb as search model.

### 3.5 Crystallisation of a shFcγRIII:hFc1 complex

15 hlgG1 derived from the serum of a myeloma patient was used to prepare Fc-fragments (hFc1) by digestion with plasmin (Deisenhofer et al., 1976). The resulting Fc-fragments were separated from the Fab-fragments by protein A chromatography. Partially digested hlgG was removed by size exclusion chromatography with MBS (2mM MOPS, 150mM NaCl, 0.02% sodium azide, pH 7.0) as running buffer. Equimolar amounts of hFc1 and  
20 shFcγRIII were mixed and diluted with MBS to a concentration of 10mg/ml. The complex was crystallised as described under conditions indicated in table 5.

## Example 4

### 25 shFcεRII (soluble human FcεRII)

The procedure was performed according to example 1 except for the indicated changes:

### 30 4.1 Cloning and Expression

FcεRII was generated by mutating the respective wild-type cDNA (Kikutani et al., 1986) and expressed according to example 2 with the mutagenous

- 33 -

primers listed in table 5. For the expression of the protein a pET23a+ vector was chosen.

#### 4.2 Refolding and purification

5 Refolding of shFcεRII was achieved as described in example 1, with the exception that prior to rapid dilution the dissolved inclusion bodies were dialysed against 6M guanidine chloride, 20mM sodium acetate, pH 4.0. shFcεRII was refolded according to example 1 with the respective refolding  
10 buffer listed in table 6. After refolding the protein solution was dialysed against PBS, concentrated 100-fold and purified by gel filtration chromatography on Superdex 75. This yielded pure shFcεRII which was dialysed against 2mM TRIS/HCl, 150mM NaCl, 0.02% sodium azide, pH 8.0, concentrated to 10mg/ml and stored at 4°C.

#### 15 Example 5

##### **shFcγRI (soluble human FcγRI)**

The procedure was performed according to example 1 except for the indicated changes:

20

#### 5.1 Cloning and Expression

shFcγRI was generated by mutating the respective wild-type cDNA (Allen & Seed, 1988) and expressed according to example 1 with the mutagenous primers listed in table 5. For the expression of the protein a pET32a+  
25 vector was chosen, which contains after the N-terminal thioredoxin a hexahistidine-tag with a C-terminal thrombin cleavage site followed by the shFcγRI in frame with the mentioned proteins and amino acid residues. For the overexpression of the fusion protein the E.coli strain BL21(DE3) containing the plasmids pUBS and pLysS (Novagen) was used.

30

The purified inclusion bodies were solubilised in 6M guanidine-HCl, 10mM β-mercaptoethanol, 50mM Tris pH8.0 and bound to a Ni-NTA column

- 34 -

(Qiagen). The elution was performed with an imidazole gradient ranging from 0 to 1M imidazole. The eluted protein was dialysed against a 1000fold volume of 150mM NaCl, 50mM Tris pH8.0, 2mM GSH, 0.5mM GSSG for 24 hours at 4°C. After concentrating the protein solution to 25% of the initial volume, thrombin was added. After 6h of incubation at 37°C the N-terminal thioredoxin and the His-tag were removed completely as verified by N-terminal sequencing. During this digestion the shFcγRI precipitated quantitatively out of solution.

## 5.2 Refolding and purification

shFcγRI was refolded according to example 1 with the respective refolding buffer listed in table 6. After the redox potential decreased to 1mM the solution was dialysed against PBS pH8.0 and concentrated.

The refolded Protein was analysed by size exclusion chromatography, which yielded a peak of the proposed monomeric receptor and non reducing SDS-PAGE which showed a major band at 30kDa.

## Example 6

### shFcεRIa (soluble human FcεRIa)

The procedure was performed according to example 1 except for the indicated changes:

## 6.1 Cloning and Expression

shFcεRI was generated by mutating the respective wild-type cDNA (Kochan et al., 1988) and expressed according to example 1 with the mutagenous primers listed in table 5. For the expression of the protein a pET23a+ vector was chosen.

**Brief description of the figures****Fig. 1: 15% reducing SDS PAGE showing the purification of sFcγRIIb**

Lane 1: Molecular weight marker. Lane 2: E. coli lysate before induction.  
5 Lane 3: E. coli lysate 1 h after induction. Lane 4: E.coli lysate 4 h after induction. Lane 5: Purified inclusion bodies of sFcγRIIb. Lane 6: Eluate of the hIgG affinity column. Lane 7: Pooled fractions of the gel filtration column.

**Fig. 2: Equilibrium gel filtration**

10 1 μg hFc solved in 10 μl equilibration buffer (10 μg sFcγRIIb/ml PBS) was applied to a size exclusion chromatography column and the absorbance of the effluent was measured (280 nm) as a function of time. The injected Fc fragment forms a complex with the sFcγRIIb in the equilibration buffer  
15 (t = 22min). The negative peak of consumed sFcγRIIb is observed at t = 26 min.

**Fig. 3: Overall structure of human sFcγRIIb**

Stereo ribbon representation of the sFcγRIIb structure. The loops supposed  
20 to be important for IgG binding are depicted in red with some of the residues within the binding site and the conserved disulfide bridge in ball and stick representation. The potential N-glycosylation sites are shown as green balls. The termini are labeled and the β-strands are numbered consecutively for the N-terminal domain in black and for the C-terminal  
25 domain in blue. The figure was created using the programs MOLSCRIPT (Kraulis, 1991) and RENDER (Merritt and Murphy, 1994).

**Fig. 4: Interdomain contacts**

The figure shows a close-up on the residues involved in the interdomain  
30 contacts of sFcγRIIb. The amino acid residues of the N-terminal domain are depicted blue and the residues of the C-terminal domain yellow. The model is covered by a 2Fo-Fc electron density contoured at 1 σ obtained from the

- 36 -

final coordinates. Hydrogen bridges between the domains are represented by white lines. The figure was created using the program MAIN (Turk, 1992).

5 **Fig. 5a: Superposition of the two FcγRIIb domains and the CH2 domain of human IgG1**

Both domains of FcγRIIb and the CH2 domain of hlgG1 were superimposed. The N-terminal domain is depicted in blue, the C-terminal domain in red and the CH2 domain of hlgG1 in green. The respective termini are labeled and  
10 the conserved disulfide bridges are depicted as thin lines.

**Fig. 5b: Structure based sequence alignment of the sFcγRIIb domains with domains of other members of the FcR family**

The upper part of the figure shows the structure based sequence alignment of the FcγRIIb and hlgG1 Fc fragment domains performed with the program  
15 GBF-3D-FIT (Lessel & Schomburg, 1994). Amino acid residues with a  $\alpha$  distance of less than 2.0 Å in the superimposed domains are masked: lilac for matching residues between the Fc fragment domains; yellow for residues in the FcγRIIb domains; and green when they can be superimposed  
20 in all four domains. The  $\beta$ -strands are indicated below this part of the alignment and are labeled consistent with Figure 3.

The lower part of the figure shows the alignment of the amino acid sequences from the other FcγRs and the homologue FcεRIa to the profile  
25 given in the upper part of the figure using routines from the GCG package (Genetics Computer Group, 1994). The upper and lower row of numbering refer to the N- and C-terminal domains of FcγRIIb. The conserved cysteins are typed in magenta and the potential glycosylation sites in blue. Identical residues within the first domain are masked orange, those in the second  
30 domain pink and green when the residues are conserved within both domains. The less conserved third domain of FcγRI is aligned between the first and the second domains. Red arrows point to residues that are involved

in side chain contacts between the first and the second domain while blue arrows depict residues that are relevant for IgG binding. The figure was produced with the program ALSCRIPT (Barton, 1993).

5     **Fig. 6: The putative binding sites of FcγRIIb**

Solid surface representations of FcγRIIb as produced with GRASP (Nicholls et al, 1991), the color coding is according to the relative surface potential from negative (red) to positive (blue). Fig. 6a shows the molecule as in Fig. 3 by a rotation of about 90° counter-clockwise around the vertical. In Fig. 6b the molecule is rotated 90° clockwise around the same axis. Both views show the putative binding regions on the C-terminal (Fig. 6a) and the N-terminal domain (Fig. 6b). The amino acid residues discussed in the text are labeled.

15     **Fig. 7: Cα-trace of the superpositioned structures of the Fcγ-receptors**  
FcγRIII red, FcγRIIa green and FcγRIIb blue. Residues important for IgG binding are shown in ball-and-stick. The N- and C-termini are labelled.

20     **Fig. 8: Overview of the FcγRIII/Fc-fragment crystal structure in ribbon representation**

The sugar residues bound to the Fc-Fragment are indicated in ball-and-stick. The FcγRIII (blue) binds in the lower hinge region between chain-B (red) and chain-A (green) of the Fc-fragment.

25     **Fig. 9: Close-up on the binding region of the FcγRIII and the Fc-fragment**  
The colour scheme is in agreement to figure 8 and residues important for complex formation are shown in ball-and-stick.

**Fig. 10a:**

30     In the upper part of figure 10a a structure based sequence alignment of the Fc-Receptor ecto-domains is shown. Conserved residues are shaded yellow and identical residues orange. The lower part of the figure shows a part of

the alignment of human antibody sequences. Residues of the human FcγRIII in contact with the Fc-fragment in the complex crystal structure are connected by lines (black for hydrophobic interaction, red for salt bridges and blue for hydrogenbridges). Residues from the Fc-receptor in contact with the A-chain of the Fc-fragment are connected with dashed lines and those in contact with the B-chain of the Fc-fragment with solid lines. Red, blue and black lines represent charged, polar and other contacts, respectively.

**Fig. 10b:**

In the upper part of figure 10b a structure based sequence alignment of the Fc-Receptor ecto-domains is shown. Conserved residues are shaded yellow and identical residues orange. Conserved residues within the less related Kir and FcA-Receptor sequences are shaded blue. The lower part of the figure shows a part of the alignment of human antibodies with the mouse IgE (mIgE) sequence. Residues of the human FcγRIII in contact with the Fc-fragment in the complex crystal structure are connected by lines (black for hydrophobic interaction, red for salt bridges and blue for hydrogenbonds). Residues from the Fc-receptor in contact with the A-chain of the Fc-fragment are connected with dashed lines and those in contact with the B-chain of the Fc-fragment with solid lines. Red, blue and black lines represent charged, polar and other contacts, respectively.

**Fig. 11 and Fig. 12:**

Fig. 11 and Fig. 12 show an alignment of the produced sFcγR, sFcεRIa and the short form of sFcεRII and the produced sFcγR and sFcεRIa without sFcεRII, respectively.



**Table 1: Crystallographic results**

The obtained preliminary crystallographic data are shown in this table.

	<b>Orthorhombic</b>	<b>Tetragonal</b>	<b>Hexagonal</b>
<b>Space group</b>	P2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub> [19]	P4 <sub>2</sub> 2 <sub>1</sub> 2 [94]	P3 [143]
<b>Unit cell dimensions</b>	a = 40.8 Å, b = 50.9 Å, c = 80.5 Å, α = 90°, β = 90°, γ = 90°	a = 85.7 Å, b = 85.7 Å, c = 63.4 Å, α = 90°, β = 90°, γ = 90°	a = 80.9 Å, b = 80.9 Å, c = 157.0 Å, α = 90°, β = 90°, γ = 90°
<b>R<sub>merge</sub></b>	5.8%	9.8%	13.6%
<b>Resolution</b>	1.7 Å	2.7 Å	3.8 Å
<b>Unique</b>	18,040	6,616	7,210
<b>Completeness</b>	89.1%	97.1%	63.0%
<b>Multiplicity</b>	3.5	4.4	1.3
<b>V<sub>m</sub>, molecules per asymmetric unit, solvent content</b>	2.09 Å <sup>3</sup> /Da, 1 mol., 41% solvent	2.91 Å <sup>3</sup> /Da, 1 mol., 58% solvent	2.97 Å <sup>3</sup> /Da, 5 mol., 59% solvent

**Table 2: Data collection statistics**

Derivative	Space Group	No. of unique reflections	Multiplicity	Resolution (Å)	Completeness (overall/last shell) (%/%)	R <sub>m</sub> (%)	No. of sites	Phasing power
NATl	P2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	18009	3.6	1.74	92.9/86.4	5.5		
NATl	P4 <sub>2</sub> 2 <sub>1</sub> 2	6615	4.5	2.70	97.1/94.3	10.1		
NATl-Baculo	P3 <sub>1</sub> 21	3545	2.5	3.0	93.0/98.9	14.4		
UOAc	P2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	7722	4.2	2.1	96.8/95.7	7.3	1	1.79
PtPy	P2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	5520	3.9	2.3	89.7/49.6	10.5	1	1.39

$$R_m = \sum |I_h - \langle I_h \rangle| / \sum \langle I_h \rangle$$

Phasing power:  $\langle F_H \rangle / E$ , where  $\langle F_H \rangle = \sum (F_H^2 / n)^{1/2}$  is the r.m.s. heavy atom structure amplitude.

$E = \sum [(F_{PHC} - F_{PH})^2 / n]^{1/2}$  is the residual lack of closure error with  $F_{PH}$  being the structure factor amplitude and  $F_{PHC} = |F_P + F_H|$  the calculated structure factor amplitude of the derivative.

**Table 3: Refinement statistics**

	Resolution range (Å)	8.0 - 1.74 Å
	No. of unique reflections ( $F > 0\sigma(F)$ )	16252
5	R factor $R_{\text{free}}$ *	19.4 27.9
	No. of atoms per asymmetric unit protein solvent	1371 150
10	Rms deviation from ideal geometry bond length (Å) bond angle (°)	0.009 2.007
15	Average B factors (Å <sup>2</sup> ) protein main chain protein side chain solvent	18.8 25.2 36.7
	Rms deviation of bonded B factors (Å <sup>2</sup> )	4.1

20 \*  $R_{\text{free}}$ : 5% of the reflections were used as a reference data set and were not included in the refinement.

**Table 4: Residues that contribute to the interdomain contact via side chains**

25	FcγRIIb	FcγRIIa	FcγRIII	FcγRI	FcεRIa
	Asn15	Asn	Ser	Ser	Arg
	Asp20	Asp	Asp	Glu	Glu
	Gln91	Gln	Gln	Gln	Gln
	His108	His	His	His	His
30	Trp110	Trp	Trp	Trp	Trp

**Table 5: Primers used for the amplification of the FcRs**

Construct	5' - Primer	3' - Primer
sFcγRI	5' - CACCCATATGGCAGTGATCTCTTT-3'	5' - AGGACTCGAGACTAGACAGGAGTTGGTA AC-3'

- 41 -

5	sFcγRIIa	5'-ACAGT <u>CATATG</u> GCAGCTCCCC-3'	5'- AAAAA <u>AGCTT</u> CAGGGCACTTGGAC-3'
	sFcγRIIb	5'- AATT <u>CCATG</u> GGGACACCTGCAGCTCCC-3'	5'- CCCAGT <u>GTCGACAGCCT</u> AATGATCCCC-3'
	sFcγRIII	5'-AAAAAA <u>CATATG</u> CGGACTGAAG-3'	5'-AAAAA <u>AGCTT</u> AACCTTGAGTGATG-3'
	sFcεRIa	5'-GATGGC <u>CATATG</u> GCAGTCCCTCAG-3'	5'- CAATGGAT <u>CCTAAA</u> ATTGTAGCCAG-3'
	sFcεRII	5'-AAAAAA <u>CATATG</u> GAGTTGCAGG-3'	5'-TGGCTGGATCCATGCTCAAG-3'

Introduced restriction sites are underlined, start- and stop-codons are depicted as bold-italics

10 Table 6: Refolding Conditions for the FcRs

	Construct	Buffer
	sFcγRI	0.1M TRIS/HCl, 1.2M arginine, 150mM NaCl, 5mM GSH, 0.5mM GSSG, 0.02% sodium azide, pH 8.0
	sFcγRIIa	0.1M TRIS/HCl, 1.4M arginine, 150mM NaCl, 2mM GSH, 0.5mM GSSG, 0.02% sodium azide, pH 8.0
	sFcγRIIb	0.1M TRIS/HCl, 1.4M arginine, 150mM NaCl, 5mM GSH, 0.5mM GSSG, 0.02% sodium azide, pH 8.0
15	sFcγRIII	0.1M TRIS/HCl, 1.0M arginine, 150mM NaCl, 2mM GSH, 0.5mM GSSG, 0.02% sodium azide, pH 8.0
	sFcεRII	0.1M TRIS/HCl, 0.8M arginine, 150mM NaCl, 5mM GSH, 0.5mM GSSG, 0.02% sodium azide, pH 8.3

Table 7: Crystallisation Conditions for the FcRs

	Construct	Condition	Space group, cell constants	Resolution
20	sFcγRIIa	26% PEG 8000, 0.2M sodium acetate/acetic acid pH 4.6, 0.02% sodium azide	C2, a=80.4Å, b=49.7Å, c=54.6Å, a=g=90°, b=128.1°	3.0Å

- 42 -

sFcγRIIb	33% PEG 2000, 0.2M sodium acetate, 0.02% sodium azide, pH5.4	P212121, a=40.8Å, b=50.9Å, c=80.5Å, a=b=g=90°	1.7Å
sFcγRIII	22% PEG 8000, 0.1M MES/TRIS pH 7.8, 0.02% sodium azide	P22121, a=36.7Å, b=60.3Å, c=85.6Å, a=b=g=90°	2.5Å
sFcγRIII: hFc1	6% PEG 8000, 0.1M MES/TRIS pH 5.6, 0.2M Na/K tartrate, 0.02% sodium azide	P6522, a=b=115.0Å, c=303.3Å, a=b=90°, g=120°	3.3Å
sFcγRIII	22% PEG 8000, 0.1M MES/TRIS pH 7.8, 0.02% sodium azide	P22121, a=36.7Å, b=60.3Å, c=85.6Å, a=b=g=90°	2.5Å

5

### References

- Ades, E.W., Phillips, D.J., Shore, S.L., Gordon, D.S., LaVia, M.F., Black, C.M., Reimer, C.B. (1976), Analysis of mononuclear cell surfaces with fluoresceinated Staphylococcal protein A complexed with IgG antibody or heat-aggregated  $\gamma$ -globulin, J. Immunol. 117, 2119.
- Allen J.M., Seed B.; "Nucleotide sequence of three cDNAs for the human high affinity Fc receptor (FcRI)"; Nucleic Acids Res. 16:11824-11824(1988).
- Amigorena, S., Bonnerot, C., Drake, J.R., Choquet, D., Hunziker, W., Guillet, J.G., Webster, P., Sautes, C., Mellman, I., Fridman, W.H. (1992), Cytoplasmic domain heterogeneity and functions of IgG Fc receptors in B lymphocytes, Science 256, 1808-1812.
- Barton, G.C. (1993), ALSCRIPT: tool to format multiple sequence alignments, Prot. Eng. 6, 37-40.
- Bazil, V. and Strominger, J.L. (1994), Metalloprotease and serine protease are involved in cleavage of CD43, CD44, and CD16 from stimulated human granulocytes, J. Immunol. 152, 1314-1322.
- Brünger, A.T., Kuriyan, J., Karplus, M. (1987), Crystallographic R factor refinement by molecular dynamics, Science 35, 458-460.
- Burmeister, W.P., Huber, A.H., Bjorkman, P.J. (1994), Crystal structure of the complex of rat neonatal Fc receptor with Fc, Nature 372, 379-383.
- Ceuppens, J.L., Baroja, M.L., van Vaeck, F., Anderson, C.L. (1988), Defect in the membrane expression of high affinity 72kD Fc $\gamma$  receptors on phagocytic cells in four healthy subjects, J. Clin. Invest. 82, 571-578.

Collaborative computational project, Number 4 (1994), The CCP4 suite: Programs for protein crystallography, *Acta crystallogr. D* 50, 760-763.

Deisenhofer, J., Jones, T.A., Huber, R., Sjodahl, J., Sjoquist, J. (1978),  
5 Crystallization, crystal structure analysis and atomic model of the complex  
formed by a human Fc fragment and fragment B of protein A from  
*Staphylococcus aureus*, *Z. Phys. Chem.* 359, 975-985.

Deisenhofer, J. (1981), Crystallographic refinement and atomic models of  
10 a human Fc fragment and its complex with fragment B of protein A from  
*Staphylococcus aureus* at 2.9- and 2.8Å resolution, *Biochemistry* 20,  
2361-2370.

Deisenhofer J., Colman PM., Huber R., Haupt H., Schwick G.;  
15 "Crystallographic structural studies of a human Fc-fragment. I. An electron-  
density map at 4Å resolution and a partial model"; Hoppe-Seyler's *Z.*  
*Physiol. Chem.* 357:435-445(1976).

Dulau, L., Cheyrou, A., Aigle, M. (1989), Directed mutagenesis using PCR,  
20 *Nucleic Acids Res.* 17, 2873.

Ellman (1959), Tissue sulfhydryl groups, *Arch. Biochem. Biophys.* 82, 79-  
77.

25 Engelhardt, W., Geerds, C., Frey, J. (1990), Distribution, inducibility and  
biological function of the cloned and expressed human  $\beta$ Fc receptor II, *Eur.*  
*J. Immunol.* 20, 1367-1377.

Engh, R.A. and Huber, R. (1991), Accurate bond and angle parameters for  
30 X-ray protein structure refinement, *Acta crystallogr.* A47, 392-400.

Fleit, H.B., Kobasiuk, C.D., Daly, C., Furie, R., Levy, P.C., Webster, R.O. (1992), A soluble form of FcγRIII is present in human serum and other body fluids and is elevated at sites of inflammation, *Blood* 79, 2721-2728.

5 Fridman, W.H., Bonnerot, C., Daeron, M., Amigorena, S., Teillaud, J.-L., Sautès, C. (1992), Structural bases of Fcγ receptor functions, *Immunol. Rev.* 125, 49-76.

10 Fridman, W.H., Teillaud, J.-L., Bouchard, C., Teillaud, C., Astier, A., Tartour, E., Galon, J., Mathiot, C., Sautès, C. (1993), Soluble Fcγ receptors, *J. Leukocyte Biol.* 54, 504-512.

15 Gabb, H.A., Jackson, R.M., Sternberg, M.J.E. (1997), Modelling protein docking using shape complementarity, electrostatics and biochemical information, *J. Mol. Biol.* 272, 106-120.

Galon, J., Bouchard, C., Fridman, W.H., Sautès, C. (1995), Ligands and biological activities of soluble Fcγ receptors, *Immunol. Lett.* 44, 175-181.

20 Genetics Compouter Group (1994), Program Manual for the Wisconsin Package Version 8, Madison, Wisconsin.

Gordon, J. et al., (1980), The molecules controlling B lymphocytes. *Immunol. Today*, 8: 339-344.

25 Grodberg, J. and Dunn, J.J. (1988), OmpT encodes the Escherichia coli outer membrane protease that cleaves T7 RNA polymerase during purification, *J. Bacteriol.* 170, 1245-1253.

30 Hogarth, P.M., Hulett, M.D., Ierino, F.L., Tate, B., Powell, M.S., Brinkworth, R.I. (1992), Identification of the immunoglobulin binding regions (IBR) of FcγRII and FcεRI, *Immunol. Rev.* 125, 21-35.

- 46 -

Homsy, J., Meyer, M., Tateno, M., Clarkson, S., Levy, J.A. (1989), The Fc and not CD4 receptor mediates antibody enhancement of HIV infection in human cells, *Science* 244, 1357-1360.

5 Hoover, R.G., Lary, C., Page, R., Travis, P., Owens, R., Flick, J., Kornbluth, J., Barlogie, B. (1995), Autoregulatory circuits in myeloma: Tumor cell cytotoxicity mediated by soluble CD16, *J. Clin. Invest.* 95, 241-247.

10 Huber, R., Deisenhofer, J., Colman, P.M., Matsushima, M. and Palm, W. (1976), Crystallographic structure studies of an IgG molecule and an Fc fragment, *Nature* 264, 415-420.

Hulett, M.D., Witort, E., Brinkworth, R.I., McKenzie, I.F.C., Hogarth, P.M. (1994), Identification of the IgG binding site of the human low affinity  
15 receptor for IgG FcγRII, *J. Biol. Chem.* 269, 15287-15293.

Hulett, M.D., Witort, E., Brinkworth, R.I., McKenzie, I.F.C., Hogarth, P.M. (1995), Multiple regions of human FcγRII (CD32) contribute to the binding of IgG, *J. Biol. Chem.* 270, 21188-21194.

20 lerino, F.L., Powell, M.S., McKenzie, I.F.C., Hogarth, P.M. (1993), Recombinant soluble human FcγRII: Production, characterization, and inhibition of the arthus reaction, *J.Exp. Med.* 178, 1617-1628.

25 Jancarik, J. and Kim, S.H. (1991), Sparse matrix sampling: A screening method for crystallization of proteins, *J. Appl. Crystallogr.* 24, 409-411.

Jones, T.A., Zou, J.-Y., Cowan, S.W., Kjeldgaard, M. (1991), Improved methods for building protein models in electron density maps and the  
30 location of errors in these models, *Acta crystallogr.* A47, 110-119.



Kikutani H., Inui S., Sato R., Barsumian E.L., Owaki H., Yamasaki K., Kaisho T., Uchibayashi N., Hardy R.R., Hirano T., Tsunasawa S., Sakiyama F., Suemura M., Kishimoto T.; "Molecular structure of human lymphocyte receptor for immunoglobulin E"; *Cell* 47(5):657-665(1986).

5

Khayat, D., Soubrane, C., Andriew, J.M., Visonneau, S., Eme, D., Tourani, J.M., Beldjord, K., Weil, M., Fernandez, E., Jaquillat, C. (1990), Changes of soluble CD16 levels in serum of HIV patients: Correlation with clinical and biological prognostic factors, *J. Infect. Dis.* 161, 430-435.

10

Kochan J., Pettine L.F., Hakimi J., Kishi K., Kinet J.P.; "Isolation of the gene coding for the alpha subunit of the human high affinity IgE receptor"; *Nucleic Acids Res.* 16:3584-3584(1988).

Simmons D., Seed B.; "The Fc-gamma receptor of natural killer cells is a phospholipid-linked membrane protein"; *Nature* 333:568-570(1988).

15

Kraulis, P.J. (1991), MOLSCRIPT: a program to produce both detailed and schematic plots of protein structures, *J. Appl. Cryst.* 24, 946-950.

20 Leslie, A.G.W. (1997), Mosflm user guide, mosflm version 5.50, MRC Laboratory of Molecular Biology, Cambridge, UK.

Lessel, U. and Schomburg, D. (1994), Similarities between protein 3-D structures, *Protein Eng.* 7, 1175-1187.

25

Littaua, R., Kurane, I. and Ennis, F.A. (1990), Human IgG Fc receptor II mediates antibody-dependent enhancement of dengue virus infection, *J. Immunol.* 144, 3183-3186.

30 Lynch, R.G., Hagen, M., Mueller, A., Sandor, M. (1995), Potential role of FcγR in early development of murine lymphoid cells: Evidence for functional

interaction between FcγR on pre-thymocytes and an alternative, non-Ig ligand on thymic stromal cells, *Immunol. Lett.* 44, 105-109.

Mathiot, C., Teillaud, J.L., Elmalek, M., Mosseri, L., Euller-Ziegler, L.,  
5 Daragon, A., Grosbois, B., Michaux, J.L., Facon, T., Bernard, J.F., Duclos,  
B., Monconduit, M., Fridman, W.H. (1993), Correlation between serum  
soluble CD16 (sCD16) levels and disease stage in patients with multiple  
myeloma, *J. Clin. Immunol.* 13, 41-48.

10 Merritt, E.A. and Murphy, M.E.P. (1994), Raster3D Version 2.0. A program  
for photorealistic molecular graphics, *Acta Cryst.* D50, 869-873.

Metzger, H. (1992A), Transmembrane signaling: The joy of aggregation, *J.*  
*Immunol.* 149, 1477-1487.

15 Metzger, H. (1992B), The receptor with high affinity for Ig E, *Immunol. Rev.*  
125, 37-48.

Müller, S. and Hoover, R.G. (1985), T cells with Fc receptors in myeloma;  
20 suppression of growth and secretion of MOPC-315 by T alpha cells, *J.*  
*Immunol.* 134, 644-7.

Nicholls, A., Sharp, K.A., Honig, B. (1991), Protein folding and association:  
insights from the interfacial and thermodynamic properties of hydrocarbons,  
25 *Proteins* 11, 281-296.

Poo, H., Kraus, J.C., Mayo-Bond, L., Todd, R.F., Petty, H.R. (1995),  
Interaction of Fcγ receptor IIIB with complement receptor type 3 in  
fibroblast transfectants: evidence from lateral diffusion and resonance  
30 energy transfer studies, *J. Mol. Biol.* 247, 597-603.

Rappaport, E.F., Cassel, D.L., Walterhouse, D.O., McKenzie, S.E., Surrey, S., Keller, M.A., Schreiber, A.D., Schwartz, E. (1993), A soluble form of the human Fc receptor FcγRIIIa: cloning, transcript analysis and detection. *Exp. Hematol.* 21, 689-696.

5

Ravanel, K., Castelle, C., Defrance, T., Wild, T.F., Charron, D., Lotteau, V., Rabourdincombe, C. (1997), Measles virus nucleocapsid protein binds to FcγRII and inhibits human B cell antibody production. *J. Exp. Med.* 186, 269-278.

10

Roman, S., Moore, J.S., Darby, C., Muller, S., Hoover, R.G. (1988), Modulation of Ig gene expression by Ig binding factors. Suppression of alpha-H chain and lambda-2-L chain mRNA accumulation in MOPC-315 by IgA-binding factor, *J. Immunology* 140, 3622-30.

15

Sarfat, D. et al., (1988), Elevation of IgE-binding factors of serum in patients with B-cell derived chronic lymphocytic leukemia. *Blood*, 71: 94-98.

20

Sauer-Eriksson, A.E., Kleywegt, G.J., Uhlen, M., Jones, T.A. (1995), Crystal structure of the C2 fragment of streptococcal protein G in complex with the Fc domain of human IgG, *Structure* 3, 265-78.

25

Small, T., et al., (1990), B-cell differentiation following autologous, conventional or T-cell depleted bone marrow transplantation: a recapitulation of normal B-cell ontogeny. *Blood*, 76: 1647-1656.

30

Sondermann, P., Huber, R., Jacob, U. (1998B), Preparation and crystallization of active soluble human FcγRIIb derived from E.coli, *Protein Structure*, submitted.

Sondermann, P., Kutscher, C., Jacob, U., Frey, J. (1998A),  
Characterization and crystallization of soluble human Fcγ recep  
tor 11 isoforms produced in insect cells, Biochemistry, submitted.

- 5    Sondermann, P., Kutscher, C., Jacob, U., Frey, J., Analysis of complexes  
of IgG and soluble human Fcγ-Receptor II produced in insect cells and its  
crystallization, submitted.

- 10    Stengelin S., Stamenkovic I., Seed B.; "Isolation of cDNAs for two distinct  
human Fc receptors by ligand affinity cloning"; EMBO J. 7:1053-  
1059(1988).

- 15    Tax, W.J.M., Willems, H.W., Reekers, P.P.M., Capel, P.J.A., Koene, R.A.P.  
(1983), Polymorphism in mitogenic effect of IgG1 monoclonal antibodies  
against T3 antigen on human T cells, Nature 304, 445-447.

- 20    Teillaud, J.L., Brunati, S., Elmalek, M., Astier, A., Nicaise, P., Moncuit, J.,  
Mathiot, C., Job-Deslandre, C., Fridman, W.H. (1990), Involvement of  
FcR + T cells and of IgG-BF in the control of myeloma cells, Mol. Immunol.  
27, 1209-17.

Turk, D. (1992), Ph.D. Thesis, TU München, Germany.

- 25    Ulvestad, E., Matre, R., Tonder, O. (1988), IgG Fc receptors in sera from  
patients with Rheumatoid Arthritis and Systemic Lupus Erythematosus,  
Scand. J. Rheumatol., Suppl. 75, 203-208.

- 30    van de Winkel, J.G.J. and Capel, P.J.A. (1993), Human IgG Fc receptor  
heterogeneity: Molecular aspects and clinical implications, Immunol. Today  
14, 215-221.

Varin, N., Sautès, C., Galinha, A., Even, J., Hogarth, P.M., Fridman, W.H. (1989), Recombinant soluble receptors for the Fc $\gamma$  portion inhibit antibody production in vitro, Eur. J. Immunol. 19, 2263-2268.

- 5     Yang, Z., Delgado, R., Xu, L., Todd, R.F., Nabel, E.G., Sanchez, A., Nabel, G.J. (1998), Distinct cellular interactions of secreted and transmembrane Ebola virus glycoproteins, Science 279, 983-984.

- 10     Zhou, M.-J., Todd, R.F., van de Winkel, J.G.J., Petty, H.R. (1993), Cocapping of the leukoadhesin molecules complement receptor type 3 and lymphocyte function-associated antigen-1 with Fc $\gamma$  receptor III on human neutrophils. Possible role of lectin-like interactions, J. Immunol. 150, 3030-3041.

### Claims

1. Recombinant soluble Fc receptor  
characterized by the absence of transmembrane domains, signal  
5 peptide and glycosylation.
2. Recombinant Fc receptor according to claim 1,  
wherein the receptor is a FcγR or a FcεR.
- 10 3. Recombinant Fc receptor according to claim 1 or 2,  
wherein the receptor is a FcγRIIb.
4. Recombinant Fc receptor according to any one of claims 1 to 3,  
wherein the receptor is of human origin.
- 15 5. Recombinant Fc receptor according to any one of claims 1 to 4,  
wherein it contains one of the amino acids as shown in one of SEQ  
ID NOs:1-6.
- 20 6. Recombinant nucleic acid containing a sequence encoding a  
recombinant Fc receptor according to any one of claims 1 to 5.
7. Recombinant nucleic acid according to claim 6,  
wherein it contains one of the sequences as shown in one of SEQ ID  
25 NOs:7-12.
8. Recombinant nucleic acid according to claim 6 or 7,  
wherein it additionally contains expression control sequences  
operably linked to the sequence encoding the recombinant Fc  
30 receptor.
9. Recombinant nucleic acid according to any one of claims 6 to 8,

wherein it is contained on a prokaryotic expression vector, preferably a pET vector.

- 5 10. Host cell characterized by the presence of a recombinant nucleic acid according to any one of claims 6 to 8.
11. Host cell according to claim 10,  
wherein it is a prokaryotic host cell, preferably an E. coli cell.
- 10 12. Process for the determination of the amount of antibodies of a certain type in the blood, plasma or serum of a patient, characterized by the use of a recombinant soluble Fc receptor according to any one of claims 1 to 5 in an immunoassay and determination of the presence of FcR-antibody complexes.
- 15 13. Process according to claim 12, wherein the immunoassay is an ELISA and preferably a sandwich assay.
- 20 14. Process according to claim 12 or 13, wherein the antibodies to be determined are IgE antibodies and the recombinant soluble receptor is a FcεR.
- 25 15. Process according to claim 14 for the determination of a predisposition or manifestation of an allergy.
- 30 16. Process according to claim 12 or 13, wherein the antibodies to be determined are IgG antibodies and the recombinant soluble receptor is a FcγR.
17. Process for the determination of the immune status of patients with chronic diseases of the immune system, wherein a Fc receptor according to any one of claims 1 to 5 is used in a competitive

immunoassay and the amount of the corresponding sFcRs in the blood, plasma or serum of a patient is determined.

- 5 18. Process according to claim 17, wherein the chronic disease is AIDS, SLE, MM or rheumatoid arthritis.
19. Use of a recombinant soluble Fc receptor according to any one of claims 1 to 5 for the screening of substances in view of their ability to act as inhibitors of the recognition and binding of antibodies to the  
10 respective cellular receptors.
20. Use according to claim 19, wherein recombinant soluble FcγRs are used and recognition and binding of IgG antibodies is of interest.
- 15 21. Pharmaceutical composition containing as active agent a recombinant soluble FcR according to any one of claims 1 to 5.
22. Pharmaceutical composition according to claim 21 for use in the treatment or prevention of autoimmune diseases, allergies or tumor  
20 diseases.
23. Pharmaceutical composition according to claim 21 or 22 for use in the treatment of AIDS, rheumatoid arthritis or multiple myeloma, containing a recombinant soluble FcγR preferably having the amino  
25 acid sequence as shown in SEQ ID NO:1.
24. Crystalline preparation of a soluble recombinant Fc receptor according to claims 1 to 5.
- 30 25. Crystalline preparation of a soluble recombinant Fc receptor / immunoglobulin complex.



26. Use of a crystalline preparation of a recombinant soluble Fc receptor according to any one of claims 1 to 5 for the generation of crystal structure data of Fc receptors.
- 5 27. Use of a crystalline preparation of a soluble recombinant Fc receptor / immunoglobulin complex for the generation of crystal structure data of receptor / Ig complexes and their respective binding sites.
- 10 28. Use of crystal structure data obtained by the use according to claims 26 or 27 for the identification and/or preparation of Fc receptor or immunoglobulin inhibitors.
- 15 29. Use of crystal structure data obtained by the use according to claim 26 or 27 for the identification and preparation of new antibody receptors.
30. Use according to any one of claims 26 to 29 in a computer-aided modelling program.
- 20 31. FcR inhibitor characterized in that it has a three-dimensional structure which is complementary to the recombinant soluble FcR according to any one of claims 1 to 5.
- 25 32. Immunoglobulin-inhibitor, characterized in that it has a three-dimensional structure which is complementary to an Fc receptor binding site of an immunoglobulin.
- 30 33. Pharmaceutical composition containing as active agent a FcR inhibitor according to claim 31.
34. Pharmaceutical composition containing as active agent an immunoglobulin inhibitor according to claim 32.

35. Pharmaceutical composition according to claim 33 or 34 for use in the treatment or prevention of diseases which are due to overreactions or faulty reactions of the immune system.
- 5 36. Pharmaceutical composition according to claim 33, 34 or 35 for the treatment or prevention of allergies, autoimmune diseases or an anaphylactic shock.
- 10 37. Use of a molecule for the modulation of the interaction between Fc receptor and immunoglobulin, characterized in that the molecule is designed or identified using crystal structure data obtained from crystalline preparations according to claims 24 or 25.
- 15 38. Use according to claim 37, wherein the modulation is partial or complete inhibition of binding between Fc receptor and immunoglobulin.
39. Fc receptor according to claims 1-5, bound to a solid phase.
- 20 40. Fc receptor according to claim 39, wherein the solid phase is a chromatography carrier material.
- 25 41. Use of a chromatography carrier material according to claim 40 for the adsorption of immunoglobulins from the blood, plasma or serum of a patient or from culture supernatants of immunoglobulin producing cells.
- 30 42. Use according to claim 41 for the enrichment of antibodies from a patient's blood, serum or plasma or from culture supernatants of immunoglobulin producing cells for the conduction of further tests.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5A




FIG. 5B






FIG. 6



FIG. 7




FIG. 8




FIG. 9

10/14





FIG. 10a



sFcγRIIa	---MAAPPKAVLKLEPP-WINVLQEDSVTLTCQGARSPESDSIQWFHN-GNLIPHTHTQPS	55
sFcγRIIb	MGTAAAPPKAVLKLPEQ-WINV LQEDSVTLTCRGTHSPESDSIQWFHN-GNLIPTHTQPS	58
sFcγRIII	-MRTEDLPKAVVFLEPQ-WYSVLEKDSVTLLKCQGAYS PEDNSTQWFHN-ESLISSQASSY	57
sFcγRI	-----MAVISLQPP-WVS VFQEETVTLHCEVLHLPGSSSTQWFLN-GTATQTSTPSY	50
sFcεRIa	---MAVPQKP KVSLNPP-WNRIFKGENVT LTCNGNNFFEV SSTKW FHN-GSLSEETNSSL	55
sFcεRII	-MELQVSSG FVCNTCPEKWINFORK-----C---YYFGKGTKQVWHARYACDDMEGQLV	50
	* * . . *	:: :*
sFcγRIIa	YRFKANND SG-EYTCQTGQTSLSDPVH LTVLSEWL V-LQTPHLEFQEGETIMLRCHSWK	113
sFcγRIIb	YRFKANND SG-EYTCQTGQTSLSDPVH LTVLSEWL V-LQTPHLEFQEGETIVLRCHSWK	116
sFcγRIII	FIDAATVN DSG-EYRCQTNLSTLSDPVQL EVHIGWLL-LQAPRVVFKEEDPIHLRCHSWK	115
sFcγRI	RITSASVN DSG-EYRCQRGLSGRSDFI QLEIHRGWLL-LQVSSRVFTEGEPLALRCHAWK	108
sFcεRIa	NIVNAKFEDSG-EYKCQHQQVNESEP VYLEVFS DWLL-LQASA EVMMEGQPLFLRCHGWR	113
sFcεRII	SIHSPEEQD FLTKHASHTGSWIGLR NLDLKG EF IWDGSHVDYSNWAPGEPTS-RSQGED	109
	. : * :: :	: * *: . . *..
sFcγRIIa	DKPLVKVTF FQNGK-SQKFSRLDPTFS IPQANHSHSGDYHCTGNIGYT LFSSKPVTITVQ	172
sFcγRIIb	DKPLVKVTF FQNGK-SKKFSRSDPNFS IPQANHSHSGDYHCTGNIGYTLYSSKPVTITVQ	175
sFcγRIII	NTALHKVTYLQNGK-DRKYFHNSDFHI PKATLKD S GSYFCRGLVGSKNVSSSETVNITIT	174
sFcγRI	DKLVYNVLYYRNGK-AFKFFHWNSNLTI LKTNI SHNGTYHCSG-MGKHRYTSAGISVTVK	166
sFcεRIa	NWDVYKVLYYK DGE-ALKY WYENHNISI TNATVEDSGTY YCTGKVWQLDYESEPLNITVI	172
sFcεRII	CVMMRGSGRW NDAFC DRKLGA WCDRLATCTPPASEGSAESMGPD SRPDPDGR LPTPSAP	169
	:	:. * : . * . *
sFcγRIIa	VP-----	174
sFcγRIIb	APSSSPMGII-----	185
sFcγRIII	QG-----	176
sFcγRI	ELFPAPVLNASVT SP LLEG NLVLTSC ET KLLLQRP GLQLYFSFYMGSKTL RGRNTS SEYQ	226
sFcεRIa	KAPREKYWLQF-----	183
sFcεRII	LHS-----	172
sFcγRIIa	-----	
sFcγRIIb	-----	
sFcγRIII	-----	
sFcγRI	ILTARRED SGLYWCEAATEDGNVLKR SPELELQVLGLQLPTPV	269
sFcεRIa	-----	
sFcεRII	-----	

13 / 14

**Alignment the produced sFcγR and sFcεR1a without sFcεR1I**

```

sFcγRIIa      ---MAAPPKAVLKLEPPWINVLQEDSVTLTCQGARSPESDSIQWFHNGNLIPTHTQPSYR  57
sFcγRIIb      MGTFAAPPKAVLKLEPQWINVLQEDSVTLTCRGTHSPESDSIQWFHNGNLIPTHTQPSYR  60
sFcγRIII      -MRTEDLPKAVVFLEPQWYSVLEKDSVTLKCQGAYSPEDNSTQWFHNESLISSQASSYFI  59
sFcγRI        -----MAVISLQPPWVSVFQEETVTLHCEVLHLPGSSSTQWFLNGTATQTSTPSYRI  52
sFcεRIa       ---MAVPQKPKVSLNPPWNRIFKGENVTLTGNGNNFFEVSSTKWPHNGSLSEETNSSLNI  57
               . : * : * : * : : : . : * * * : * . : * : * * : . :

sFcγRIIa      FKANNNDSGEYTCQTGQTSLSDPVHLTVLSEWLVLQTPHLEFQEGETIMLRCHSWKDKPL  117
sFcγRIIb      FKANNNDSGEYTCQTGQTSLSDPVHLTVLSEWLVLQTPHLEFQEGETIVLRCHSWKDKPL  120
sFcγRIII      DAATVNDSGEYRCQTNLSTLSDPVQLEVHIGWLLQAPRWVFKEEDPIHLRCHSWKNTAL  119
sFcγRI        TSASVNDSGEYRCQRGSLGRSDPIQLEIHRGWLLQVSSRVFTEGEPLALRCHAWKDKLV  112
sFcεRIa       VNAKFEDSGEYKQHQVNESEPVYLEVFSDWLLQASAEVVMEGQPLFLRCHGWRNWDV  117
               * . : * * * * * * * : * : * : * : * : * : * : * : * : * :

sFcγRIIa      VKVTFQNGKSKFQSRDLPTFSIPQANHSHSGDYHCTGNIGYTLFSSKPVTTITVQVP---  174
sFcγRIIb      VKVTFQNGKSKKFSRSDPNFSIPQANHSHSGDYHCTGNIGYTLYSSKPVTTITVQAPSSS  180
sFcγRIII      HKVTYLQNGKDRKYFHHNSDFHIPKATLKDSGSYFCRGLVGSKNVSETVNITITQG---  176
sFcγRI        YNVLYYRNGKAFKFFHWNNSLTILKTNISHNGTYHCSG-MGKHRYTSAGISVTVKELFPA  171
sFcεRIa       YKVIYYKDGEALKYWYENHNISITNATVEDSGTYCYCTGKVVQLDYESEPLNITVIKAPRE  177
               : * : : * : * : : : * : : . . . * * * * : * : : * :

sFcγRIIa      -----
sFcγRIIb      PMGII-----  185
sFcγRIII      -----
sFcγRI        PVLNASVTSPLLEGNLVTLSCETKLLLQRPGLQLYFSFYMGSKTLRGRNTSSEYQILTAR  231
sFcεRIa       KYWLQF-----  183

sFcγRIIa      -----
sFcγRIIb      -----
sFcγRIII      -----
sFcγRI        REDSGLYWCEAATEDGNVLKRSPELELQVLGLQLPTPV  269
sFcεRIa       -----

```

FIG. 12



**Protein sequences of the examples (mutated to yield soluble forms)**

&gt;sFcγRI

SEQ ID NO: 1

MAVISLQPPWVSVFQEETVTLHCEVLHLPGSSSTQWFLNGTATQTSTPSYRITSASVND  
GEYRCQRGLSGRSDPIQLEIHRGWLLQVSSRVFTEGEPLALRCHAWKDKLVYNVLYYRN  
GKAFKFFHWNSNLTILKTNISHNGTYHCSGMGKHRYTSAGISVTVKELFPAPVLNASVTS  
PLLEGNLVTLSCEKLLLRPGLQLYFSFYMGSKTLRGRNTSSEYQILTARREDSGLYWC  
EAATEDGNVLKRSPELELQVLGLQLPTPV

SEQ ID NO: 2

&gt;sFcγRIIa

MAAPPKAVLKLEPPWINVLQEDSVTLTCQGARSPESDSIQWFHNGNLIPTHTQPSYRFKA  
NNNDSGEYTCQTGQTSLSDPVHLTVLSEWLVLQTPHLEFQEGETIMLRCHSWKDKPLVKV  
TFFQNGKSKQFSRLDPTFSIPQANHSHSGDYHCTGNIGYTLFSSKPVTTITVQVP

SEQ ID NO: 3

&gt;sFcγRIIb

AGTPAAPPKAVLKLEPPQWINVLQEDSVTLTCRGTSHSPESDSIQWFHNGNLIPTHTQPSYR  
FKANNNDSGEYTCQTGQTSLSDPVHLTVLSEWLVLQTPHLEFQEGETIVLRCHSWKDKPL  
VKVTFFQNGKSKKFSRSDPNFSIPQANHSHSGDYHCTGNIGYTLYSSKPVTTITVQAPSSS  
PMGII

SEQ ID NO: 4

&gt;sFcγRIII

MRTEDLPKAVVFLEPQWYSVLEKDSVTLKCQGAYPEDNSTQWFHNESLISSQASSYFID  
AATVNDSGEYRCQTNLSTLSDPVQLEVHIGWLLQAPRWVFKEEDPIHLRCHSWKNTALH  
KVTYLQNGKDRKYFHHNSDFHIPKATLKDSGSYFCRGLVGSKNVSSSETVNITITQG

SEQ ID NO: 5

&gt;sFcεRIa

MAVPQKPKVSLNPPWNRIFKGENVTLCNGNFFEVSSSTKWFHNGSLSEETNSSLNIVN  
AKFEDSGEYKQHQVNESEPVYLEVFSDWLLQASAEVVMEGQPLFLRCHGWRNWDVYK  
VIYYKDGEALKYWYENHNISITNATVEDSGTYCYCTGKVVQLDYESEPLNITVIKAPREKY  
W'LQF

SEQ ID NO: 6

&gt;sFcεRII

MDTTQSLKQLEERAARNVSQVSKNLESHHGDQMTQKSQSTQISQELEELRAEQQRLKSQD  
LELSWNLNGLQADLSSFKSQELNERNEASDLLERLREEVTKLRMELQVSSGFVCNTCPEK  
WINFQRKCYFYGKGTKQWVHARYACDDMEGQLVSIHSPEEQDFLTKHASHTGSWIGLRNL  
DLKGEFIWVDGSHVDYSNWPAGEPTSRSQGEDCVMMRGSGRWNDAFCDRKLGAWVCDRLA  
TCTPPASEGSAESMGPDSPDPDGRLPTPSAPLHS

**DNA sequences of the examples (mutated to yield soluble forms)**

SEQ ID NO: 7

>sFyRI

```

1 CATATGGCAG TGATCTCTTT GCAGCTCCA TGGGTCAGCG TGTTCCAAGA GGAAACCGTA 60
61 ACCTTGCACT GTGAGGTGCT CCATCTGCCT GGGAGCAGCT CTACACAGTG GTTTCTCAAT 120
121 GGCACAGCCA CTCAGACCTC GACCCCCAGC TACAGAATCA CCTCTGCCAG TGTCATGAC 180
181 AGTGGTGAAT ACAGGTGCCA GAGAGGTCTC TCAGGGCGAA GTGACCCCAT ACAGCTGGAA 240
241 ATCCACAGAG GCTGGCTACT ACTGCAGGTC TCCAGCAGAG TCTTCACGGA AGGAGAACCT 300
301 CTGGCCTTGA GGTGTCATGC GTGGAAGGAT AAGCTGGTGT ACAATGTGCT TTAATATCGA 360
361 AATGGCAAAG CCTTTAAGTT TTTCCACTGG AATTCTAACC TCACCATTCT GAAAACCAAC 420
421 ATAAGTCACA ATGGCACCTA CCATTGCTCA GGCATGGGAA AGCATCGCTA CACATCAGCA 480
481 GGAATATCTG TCACTGTGAA AGAGCTATTT CCAGCTCCAG TGCTGAATGC ATCTGTGACA 540
541 TCCCCACTCC TGGAGGGGAA TCTGGTCACC CTGAGCTGTG AAACAAAGTT GCTCTTGCAG 600
601 AGGCCTGGTT TGCAGCTTTA CTTCTCCTTC TACATGGGCA GCAAGACCCCT GCGAGGCAGG 660
661 AACACACCTT CTGAATACCA AATACTAACT GCTAGAAGAG AAGACTCTGG GTTATACTGG 720
721 TGGCAGGCTG CCACAGAGGA TGGAAATGTC CTTAAGCGCA GCCCTGAGTT GGAGCTTCAA 780
781 GTGCTTGCC TCCAGTTACC AACTCCTGTC TAGTCTCGAG 820

```

SEQ ID NO: 8

>sFcyRIIa

```

1 CATATGGCAG CTCCCCAAA GGCTGTGCTG AAACCTGAGC CCCCCTGGAT CAACGTGCTC 60
61 CAGGAGGACT CTGTGACTCT GACATGCCAG GGGGCTCGCA GCCCTGAGAG CGACTCCATT 120
121 CAGTGGTTCC ACAATGGGAA TCTCATTTCC ACCCACACGC AGCCCAGCTA CAGGTTCAAG 180
181 GCCAACAACA ATGACAGCGG GGAGTACACG TGCCAGACTG GCCAGACCAG CCTCAGCGAC 240
241 CCTGTGCATC TGAATGTGCT TTCCGAATGG CTGGTGCTCC AGACCCCTCA CCTGGAGTTC 300
301 CAGGAGGGAG AAACCATCAT GCTGAGGTGC CACAGCTGGA AGGACAAGCC TCTGGTCAAG 360
361 GTCACATTCT TCCAGAATGG AAAATCCCAG AAATTCTCCC GTTTGGATCC CACCTTCTCC 420
421 ATCCCACAAG CAAACCACAG TCACAGTGGT GATTACCACT GCACAGGAAA CATAGGCTAC 480
481 ACGCTGTCTT CATCCAAGCC TGTGACCATC ACTGTCCAAG TGCCCTGAAG CTT 533

```

SEQ ID NO: 9

>sFcyRIIb

```

1 CCATGGGGAC ACCTGCAGCT CCCCCAAAGG CTGTGCTGAA ACTCGAGCCC CAGTGGATCA 60
61 ACGTGCTCCA GGAGGACTCT GTGACTCTGA CATGCCGGGG GACTCACAGC CCTGAGAGCG 120
121 ACTCCATTCA GTGGTTCCAC AATGGGAATC TCATTCCCAC CCACACGCAG CCCAGCTACA 180
181 GGTTCAGGGC CAACAACAAT GACAGCGGGG AGTACACGTG CCAGACTGGC CAGACCAGCC 240
241 TCAGCGACCC TGTGCATCTG ACTGTGCTTT CTGAGTGGCT GGTGTCCAG ACCCTCACC 300
301 TGGAGTTCCA GGAGGGAGAA ACCATCGTGC TGAGGTGCCA CAGCTGGAAG GACAAGCCTC 360
361 TGGTCAAGGT CACATTCTTC CAGAATGGAA AATCCAAGAA ATTTTCCCGT TCGGATCCCA 420
421 ACTTCTCCAT CCCACAAGCA AACCACAGTC ACAGTGGTGA TTACCACTGC ACAGGAAACA 480
481 TAGGCTACAC GCTGTACTCA TCCAAGCCTG TGACCATCAC TGTCCAAGCT CCCAGCTCTT 540
541 CACCGATGGG GATCATTTAG GCTGTCGAC 569

```

SEQ ID NO: 10

>sFcyRIII

```

1 CATATGCGGA CTGAAGATCT CCCAAAGGCT GTGGTGTTC TGGAGCCTCA ATGGTACAGC 60
61 GTGCTTGAGA AGGACAGTGT GACTCTGAAG TGCCAGGGAG CCTACTCCCC TGAGGACAAT 120
121 TCCACACAGT GGTTCACAA TGAGAGCCTC ATCTCAAGCC AGGCCTCGAG CTACTTCATT 180
181 GACGCTGCCA CAGTCAACGA CAGTGGAGAG TACAGGTGCC AGACAAACCT CTCCACCCTC 240
241 AGTGACCCGG TGCAGCTAGA AGTCCATATC GGTGCTGCTG TGCTCCAGGC CCCTCGGTGG 300
301 GTGTTCAAGG AGGAAGACCC TATTCACCTG AGGTGTCACA GCTGGAAGAA CACTGCTCTG 360
361 CATAAGGTCA CATATTTACA GAATGGCAAA GACAGGAAGT ATTTTCATCA TAATTCTGAC 420
421 TTCCACATTC CAAAAGCCAC ACTCAAAGAT AGCGGCTCCT ACTTCTGCAG GGGGCTTGTT 480
481 GGGAGTAAAA ATGTGTCTTC AGAGACTGTG AACATCACCA TCACTCAAGG TTAAGCTT 538

```

SEQ ID NO: 11

>sFceRIa

```

1 CATATGGCAG TCCCTCAGAA ACCTAAGGTC TCCTTGAACC CTCCATGGAA TAGAATATTT 60
61 AAAGGAGAGA ATGTGACTCT TACATGTAAT GGGAAACAATT TCTTTGAAGT CAGTTCACC 120
121 AAATGGTTCC ACAATGGCAG CCTTTCAGAA GAGACAAATT CAAGTTTGAA TATTGTGAAT 180
181 GCCAAATTTG AAGACAGTGG AGAATACAAA TGTCAGCACC AACAGTTAA TGAGAGTGAA 240
241 CCTGTGTACC TGGAAGTCTT CAGTGAAGTG CTGCTCCTTC AGGCCTCTGC TGAGGTGGTG 300
301 ATGGAGGGCC AGCCCTCTTT CCTCAGGTGC CATGGTTGGA GGAACCTGGG TGTTGTACAAG 360
361 GTGATCTATT ATAAGGATGG TGAAGCTCTC AAGTACTGGT ATGAGAACCA CAACATCTCC 420
421 ATTACAAATG CCACAGTTGA AGACAGTGGA ACCTACTACT GTACGGGCAA AGTGTGGCAG 480
481 CTGGACTATG AGTCTGAGCC CCTCAACATT ACTGTAATAA AAGCTCCGCG TGAGAAGTAC 540
541 TGGCTACAAT TTTAGGATCC 560

```

## SEQ ID NO: 12

&gt;sFceRII

```

1  CATATGGAGT TGCAGGTGTC CAGCGGCTTT GTGTGCAACA CGTGCCCTGA AAAGTGGATC 60
61  AATTTCCAAC GGAAGTGCTA CTA CTTCGGC AAGGGCACCA AGCAGTGGGT CCACGCCCGG 120
121 TATGCCTGTG ACGACATGGA AGGGCAGCTG GTCAGCATCC ACAGCCCGGA GGAGCAGGAC 180
181 TTCCTGACCA AGCATGCCAG CCACACCGGC TCCTGGATTG GCCTTCGGAA CTTGGACCTG 240
241 AAGGGGGAGT TTATCTGGGT GGATGGGAGC CACGTGGACT ACAGCAACTG GGCTCCAGGG 300
301 GAGCCCACCA GCCGGAGCCA GGGCGAGGAC TGC GTATGA TGCGGGGCTC CGGTCGCTGG 360
361 AACGACGCCT TCTGCGACCG TAAGCTGGGC GCCTGGGTGT GCGACCGGCT GGCCACATGC 420
421 ACGCCGCCAG CCAGCGAAGG TTCCGCGGAG TCCATGGGAC CTGATTCAAG ACCAGACCTT 480
481 GACGGCCGCC TGCCACCCC CTCTGCCCCC CTCCACTCTT GAGCATGGAT CC 532

```

## SEQ ID NO:13

human FcγRIIb2

```

1  ggctgtgact gctgtgctct gggcgccact cgctccaggg agtgatggga atcctgtcat
61  ttttacctgt ccttgccact gagagtgact gggctgactg caagtccccc cagccttggg
121  gtcatatgct tctgtggaca gctgtgctat tcttggtccc tgttgctggg acacctgcag
181  ctccccaaaa ggctgtgctg aaactcgagc cccagtggat caacgtgctc caggaggact
241  ctgtgactct gacatgccgg gggactcaca gccctgagag cgactccatt cagtggttcc
301  acaatgggaa tctcattccc acccacagc agcccagcta caggttcaag gccaacaaca
361  atgacagcgg ggagtacacg tgcagactg gccagaccag cctcagcgac cctgtgcac
421  tgacagtgtc ttctgagtggt ctggtgctcc agaccctca cctggagtcc caggagggag
481  aaaccatcgt gctgaggtgc cacagctgga aggacaagcc tctggtcaag gtcacattct
541  tccagaatgg aaaatccaag aaattttccc gttcggatcc caacttctcc atcccacaag
601  caaaccacag tcacagtggg gattaccatt gcacaggaaa cataggctac acgtgtact
661  catccaagcc tgtgaccatc actgtccaag ctcccagctc ttcaccgatg gggatcattg
721  tggctgtggg cactgggatt gctgtagctg ccattgttgc tgctgtagtg gccttgatct
781  actgcaggaa aaagcggatt tcagccaatc cactaatcc tgatgagggt gacaaagttg
841  gggctgagaa cacaatcacc tattcacttc tcatgcaccc ggatgctctg gaagagcctg
901  atgaccagaa ccgtatttag tctccattgt cttgcattgg gatttgagaa gaaatcagag
961  agggaagatc tggatatttc tggcctaaat tccccttggg gaggacaggg agatgctgca
1021 gttccaaaag agaaggttcc tccagagtc atctacctga gtctgaagc tccctgtcct
1081 gaaagccaca gacaatatgg tcccaaatgc ccgactgcac cttctgtgct tcagctcttc
1141 ttgacatcaa ggctcttccg ttcacatcc acacagcaa tccaattaat caaaccactg
1201 ttattaacag ataatagcaa cttgggaaat gcttatgtta caggttacgt gagaacaatc
1261 atgtaaatct atatgatttc agaaatgtta aaatagacta acctctacca gcacattaac
1321 agtgattgtt tctgggtgat aaaattattg atgattttta ttttctttat ttttctataa
1381 agatcatata ttacttttat aataaaacat tataaaaac

```

## SEQ ID NO:14

human FcεRIα

```
1      agatctcagc acagtaagca ccaggagtcc atgaagaaga tggctcctgc catggaatcc
61     cctactctac tgtgtgtagc cttactgttc ttcgctccag atggcgtgtt agcagtccct
121    cagaaaccta aggtctcctt gaaccctcca tggaatagaa tatttaaagg agagaatgtg
181    actcttacat gtaatgggaa caatttcttt gaagtcagtt ccaccaaagt gttccacaat
241    ggcagccttt cagaagagac aaattcaagt ttgaatattg tgaatgccaa atttgaagac
301    agtggagaat acaaatgtca gcaccaacaa gttaatgaga gtgaacctgt gtacctggaa
361    gtcttcagtg actggctgct ccttcaggcc tctgctgagg tggatgatga gggccagccc
421    ctcttcctca ggtgccatgg ttggaggaac tgggatgtgt acaaggtgat ctattataag
481    gatggtgaag ctctcaagta ctggtatgag aaccacaaca tctccattac aaatgccaca
541    gttgaagaca gtggaaccta ctactgtacg ggcaaagtgt ggcagctgga ctatgagtct
601    gagccccctc acattactgt aataaaagct ccgcgtgaga agtactggct acaatttttt
661    atcccattgt tgggtggtgat tctgtttgct gtggacacag gattatztat ctcaactcag
721    cagcaggtca catttctctt gaagattaag agaaccagga aaggcttcag acttctgaac
781    ccacatccta agccaaaccc caaaaacaac tgatataatt aactcaagaa atatttgcaa
841    cattagtttt tttccagcat cagcaattgc tactcaattg tcaaacacag cttgcaatat
901    acatagaaac gtctgtgctc aaggatttat agaaatgctt cattaaactg agtgaaactg
961    attaagtggc atgtaatagt aagtgtctca ttaacattgg ttgaataaat gagagaatga
1021   atagattcat ttattagcat ttgtaaaaga gatgttcaat ttagatct
```

## SEQ ID NO:15

human mRNA for high affinity Fc receptor (FcγRI)

```
1      gacagatttc actgctccca ccagcttgga gacaacatgt ggttcttgac aactctgctc
61     ctttgggttc cagttgatgg gcaagtggac accacaaagg cagtgatctc ttgacagcct
121    ccatgggtca gcgtgttcca agaggaaacc gtaaccttgc actgtgaggt gctccatctg
181    cctgggagca gctctacaca gtggtttctc aatggcacag ccaactcagac ctgaccccc
241    agctacagaa tcacctctgc cagtgatcaat gacagtgggtg aatacagggtg ccagagaggt
301    ctctcagggc gaagtgacct catacagctg gaaatccaca gaggttggt actactgcag
361    gtctccagca gagtcttcac ggaaggagaa cctctggcct tgagggtgtca tgcgtggaag
421    gataagctgg tgtacaatgt gctttactat cgaaatggca aagcctttaa gtttttccac
481    tggaattcta acctcaccat tctgaaaacc aacataagtc acaatggcac ctaccattgc
541    tcaggcatgg gaaagcatcg ctacacatca gcaggaatat ctgtcactgt gaaagagcta
601    tttccagctc cagtgtgaa tgcactctgtg acatccccac tcctggaggg gaatctggtc
661    accctgagct gtgaaacaaa gttgctcttg cagaggcctg gtttgagct ttacttctcc
721    ttctacatgg gcagcaagac cctgcgaggc aggaacacat cctctgaata ccaaatacta
781    actgctagaa gagaagactc tgggttatac tgggtcgagg ctgccacaga ggatggaaat
841    gtctttaagc gcagccctga gttggagctt caagtgttg gcctccagtt accaactcct
901    gtctggtttc atgtcctttt ctatctggca gtgggaataa tgtttttagt gaacactgtt
961    ctctgggtga caatacgtaa agaactgaaa agaaagaaaa agtgggattt agaaatctct
1021   ttggattctg gtcattgagaa gaaggtaact tccagccttc aagaagacag acatttagaa
1081   gaagagctga aatgtcagga acaaaaagaa gaacagctgc aggaaggggt gcaccggaag
1141   gagccccagg gggccacgta gcagcggctc agtgggtggc catcgatctg gaccgtcccc
1201   tgcccacttg ctccccgtga gcaactgcgt caaacatcca aaagttcaac aacaccagaa
1261   ctgtgtgtct catggtatgt aactcttaaa gcaaataaat gaactgactt caaaaaaaaa
1321   a
```

## SEQ ID NO:16

human FcγRIIa

```
1      cccaaatgtc tcagaatgta tgtcccagaa acctgtggct gcttcaacca ttgacagttt
61     tgctgctgct ggcttctgca gacagtcaag ctgcagctcc cccaaaggct gtgctgaaac
121    ttgagcccc gtggatcaac gtgctccagg aggactctgt gactctgaca tgccaggggg
181    ctgcagccc tgagagcgac tccattcagt ggttccacaa tgggaatctc attcccaccc
241    acacgcagcc cagctacagg ttcaaggcca acaacaatga cagcggggag tacacgtgcc
301    agactggcca gaccagcctc agcgacctg tgcactctgac tgtgctttcc gaatggctgg
361    tgctccagac cctcacctg gagttccagg agggagaaac catcatgctg aggtgccaca
421    gctggaagga caagcctctg gtcaaggcca cattcttcca gaatggaaaa tcccagaaat
481    tctcccgttt ggatcccacc ttctccatcc cacaagcaaa ccacagtcac agtgggtgatt
541    accactgcac aggaacata ggctacacgc tgttctcatc caagcctgtg accatcactg
601    tccaagtgcc cagcatgggc agctcttcac caatggggat cattgtggct gtggtcattg
661    cgactgctgt agcagccatt gttgctgctg tagtggcctt gatctactgc aggaaaaagc
721    ggatttcagc caattccact gatcctgtga aggctgccca atttgagcca cctggacgtc
781    aaatgattgc catcagaaag agacaacttg aagaaaccaa caatgactat gaaacagctg
841    acggcggtca catgactctg aaccccaggg cacctactga cgatgataaa aacatctacc
901    tgactcttcc tcccacgac catgtcaaca gtaataacta aagagtaacg ttatgccatg
961    tggtcatact ctgagcttgc tgatggatga caaaaagagg ggaattgtta aaggaaaatt
1021   taaatggaga ctggaaaaat cctgagcaaa caaaaccacc tggcccttag aaatagcttt
1081   aactttgctt aaactacaaa cacaagcaaa acttcacggg gtcatactac atacaagcat
1141   aagcaaaact taacttggat ctttctgggt aaatgcttat gttagaaata agacaacccc
1201   agccaatcac aagcagccta ctaacatata attaggtgac tagggacttt ctaagaagat
1261   acctaccccc aaaaaacaat tatgtaattg aaaaccaacc gattgccttt attttgcttc
1321   cacattttcc caataaatac ttgctgtgta ctttttgcca ctggaacact aaacttcatt
1381   aattgcgcct cagatttttc cttaacatc tttttttttt ttgacagagt ctcaatctgt
1441   taccaggctt ggagtgcagt ggtgctatct tggtcactg caaacccgcc tcccaggttt
1501   aagcgattct tatgcctcag cctccagta gctgggatta gaggcattgt ccatcatacc
1561   cagctaattt ttgtattttt ttttttttat ttttagtaga gacagggttt cgcaatggtg
1621   gccaggccga tctcgaactt ctggcctcta gcatctgcc cgcctcggcc tcccaaagtg
1681   ctgggatgac cgcctcagcc ccaatgtcca gcctctttaa catcttcttt cctatgcctt
1741   ctctgtggat ccctactgct ggtttctgcc ttctccatgc tgagaacaaa atcacctatt
1801   cactgcttat gcagtgcgaa gctccagaag acaaaagagc ccaattacca gaaccacatt
1861   aagtctccat tgttttgctt tgggatttga gaagagaatt agagaggtga ggatctggta
1921   tttcctggac taaattccct tggggaagac gaagggatgc tgcagttcca aaagagaagg
1981   actcttccag agtcatctac ctgagtccca aagctccctg tctgaaagc cacagacaat
2041   atggtcccaa atgactgact gcacctctct tgctcagcc gttcttgaca tcaagaatct
2101   tctgttccac atccacacag ccaatacaat tagtcaaacc actgttatta acagatgtag
2161   caacatgaga aacgcttatg ttacaggtta catgagagca atcatgtaag tctatatgac
2221   ttcagaaatg ttaaaataga ctaacctcta acaacaaatt aaaagtgatt gtttcaaggt
2281   gatgcaatta ttgatgacct atttattttt tctataatga tcatatatta cctttgtaat
2341   aaaacattat aacaaaaac
```

## SEQ ID NO:17

human FcγRIII

```
1      tctttggtga cttgtccact ccagtgtggc atcatgtggc agctgtcct cccaactgct
61     ctgtacttc tagtttcagc tggcatgcg actgaagatc tcccaaaggc tgtggtgttc
121    ctggagcctc aatggtacag cgtgcttgag aaggacagtg tgactctgaa gtgccaggga
181    gcctactccc ctgaggacaa ttccacacag tggtttcaca atgagagcct catctcaagc
241    caggcctcga gctacttcat tgacgtgcc acagtcaacg acagtggaga gtacagggtc
301    cagacaaacc tctccaccct cagtgacccg gtgcagctag aagtccatat cggctggctg
361    ttgctccagg cccctcgggtg ggtgttcaag gaggaagacc ctattcacct gaggtgtcac
421    agctggaaga aactgtctct gcataaggtc acatatttac agaatggcaa agacaggaag
481    tattttcatc ataattctga ctccacatt ccaaaagcca cactcaaaga tagcggctcc
541    tacttctgca gggggcttgt tgggagtaaa aatgtgtctt cagagactgt gaacatcacc
601    atcactcaag gtttggcagt gtcaaccatc tcatcattct ctccacctgg gtaccaagtc
661    tctttctgct tggatgatgt actccttttt gcagtggaca caggactata tttctctgtg
721    aagacaaaca tttgaagctc aacaagagac tgggaaggacc ataaacttaa atggagaaag
781    gaccctcaag acaaatgacc cccatcccat gggagtaata agagcagtgg cagcagcatc
841    tctgaacatt tctctggatt tgcaaccca tcctctcag gcctctc
```

## SEQ ID NO: 18

## human FcεRII

1 ctctctgctta aacctctgtc tctgacggtc cctgccaatc gctctgggtcg  
accccaacac  
61 actaggagga cagacacagg ctccaaactc cactaagtga ccagagctgt gattgtgccc  
121 gctgagtgga ctgcgttgtc agggagtggag tgctccatca tcgggagaat ccaagcagga  
181 ccgccatgga ggaagggtcaa tattcagaga tcgaggagct tcccaggagg cgggtgttgca  
241 ggcgtggggac tcagatcgtg ctgctggggc tggtgaccgc cgctctgtgg gctgggctgc  
301 tgactctgct tctcctgtgg cactgggaca ccacacagag tctaaaacag ctggaagaga  
361 gggctgcccc gaacgtctct caagtttcca agaacttggg aagccaccac ggtgaccaga  
421 tggcgcagaa atcccagtc acgcagattt cacaggaact ggaggaactt cgagctgaac  
481 agcagagatt gaaatctcag gacttggagc tgtcctggaa cctgaacggg cttcaagcag  
541 atctgagcag cttcaagtcc caggaattga acgagaggaa cgaagcttca gatttgcctg  
601 aaagactccg ggaggagggtg acaaagctaa ggatggagtt gcaggtgtcc agcggccttg  
661 tgtgcaacac gtgccctgaa aagtggatca atttccaacg gaagtgtctac tacttcggca  
721 agggcaccaa gcagtgggtc cagcggcggt atgcctgtga cgacatggaa gggcagctgg  
781 tcagcatcca cagcccgag gagcaggact tctgaccaa gcatgccagc cacaccggct  
841 cctggattgg ccttcggaac ttggacctga agggagagtt tatctgggtg gatgggagcc  
901 atgtggacta cagcaactgg gctccagggg agcccaccag ccggagccag ggcgaggact  
961 gcgtgatgat gcggggctcc ggtcgtgga acgacgcctt ctgcgaccgt aagctggggc  
1021 cctgggtgtg cgaccggctg gccacatgca cgccgccagc cagcgaaggt tccgcgaggt  
1081 ccatgggacc tgattcaaga ccagaccctg acggccgcct gcccaccccc tctgcccctc  
1141 tccactcttg agcatggata cagccaggcc cagagcaaga cctgaagac cccaaccac  
1201 ggcctaaaag cctctttgtg gctgaaaggt ccctgtgaca ttttctgcca cccaaacgga  
1261 ggcagctgac acatctcccg ctctctatg gccctgcct tcccaggagt acaccccaac  
1321 agcaccctct ccagatggga gtgccccaa cagcaccctc tccagatgag agtacacccc  
1381 aacagcacc cctccagatg cagccccatc tctcagcac cccaggacct gagtatcccc  
1441 agctcagggt gtgagtcctc ctgtccagcc tgcataata aaatggggca gtgatggcct  
1501 ccc



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 99/09440

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/705 C12N1/21 C12N15/70 G01N33/53  
G01N33/68 A61K38/17 C07K17/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FR 2 739 560 A (ROUSSEL UCLAF) 11 April 1997 (1997-04-11)	1-23
Y	page 3, last line -page 6, line 28	24, 26, 28-30
X	EP 0 614 978 A (ROUSSEL-UCLAF) 14 September 1994 (1994-09-14)	1, 2, 4-13, 16, 21-23
Y	page 3, line 1 -page 7, line 33; examples --- -/--	24, 26, 28-30

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 April 2000

Date of mailing of the international search report

12/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Montero Lopez, B

## INTERNATIONAL SEARCH REPORT

Ir. ational Application No

PCT/EP 99/09440

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 623 053 A (LOUIS N. GASTINEL ET AL.) 22 April 1997 (1997-04-22)	1,4,6, 8-13, 39-42
Y	column 5, line 60 -column 8, line 67; examples	5,7,24, 26,28-30
X	WO 96 40199 A (UNIVERSITY OF PENNSYLVANIA) 19 December 1996 (1996-12-19)	1,2,4,6, 8-16, 19-22
Y	page 22, line 24 -page 24, line 14; claims 13-16,36-39; examples I,II	5,7,24, 26,28-30
X	WO 95 09002 A (UNIVERSITY OF PENNSYLVANIA) 6 April 1995 (1995-04-06)	1,2,4,6, 8-13,16, 21,39,40
Y	page 15, line 4 -page 16, line 22; claims 28-31; examples I,II	5,7,24, 26,28-30
X	EP 0 791 653 A (SCHERING BIOTECH CORPORATION) 27 August 1997 (1997-08-27)	1,2,4,6, 8-11,21, 22
Y	column 5, line 56 -column 10, line 1; example III	5,7,24, 26,28-30
X	JÉRÔME GALON ET AL.: "Ligands and biological activities of soluble Fcgamma receptors" IMMUNOLOGY LETTERS, vol. 44, January 1995 (1995-01), pages 175-181, XP000574018	1-4,6, 8-13,15, 16,19-22
Y	page 175, right-hand column, last paragraph -page 177, left-hand column, paragraph 1 page 179, right-hand column, paragraph 3 -page 180, right-hand column, paragraph 3	5,7,24, 26,28-30
X	EP 0 321 842 A (KISHIMOTO, TADAMITSU ) 28 June 1989 (1989-06-28)	1,2, 4-19,21, 22
Y	column 2, line 10 -column 9, line 43; figure 4; examples B,,1-3,6,,7	24,26, 28,30
	--- -/--	

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/09440

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 259 615 A (KISHIMOTO, TADAMITSU) 16 March 1988 (1988-03-16)	1,2, 4-15, 17-19, 21,22 24,26, 28-30
Y	page 2, line 1 -page 37, line 13 -----	
Y	JANET M. ALLEN ET AL.: "Isolation and expression of functional high-affinity Fc receptor complementary DNAs" JOURNAL OF CRYSTAL GROWTH, vol. 93, no. 1-4, November 1988 (1988-11) - December 1988 (1988-12), pages 378-381, XP000002750 Amsterdam, NL abstract; figure 4 -----	5,7
Y	WILHELM P. BURMEISTER ET AL.: "Crystal structure at 2.2A resolution of the MHC-related neonatal Fc receptor" NATURE, vol. 372, no. 6504, 24 November 1994 (1994-11-24), pages 336-343, XP002135452 LONDON GB the whole document -----	24,26, 28-30
P,X	WO 99 05271 A (GOULD, HANNAH ET AL.) 4 February 1999 (1999-02-04)  page 4, line 6 - line 19 page 5, line 22 -page 6, line 2 page 7, line 22 -page 8, line 23 page 10, line 31 -page 11, line 31 -----	1,2,4,6, 8-15,21, 22,39

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/ 09440

### Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 31-38  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
See FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

#### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/EP 99 09440

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 31-38

Present claims 31-38 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is not to be found, however, for any specific example of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, no search has been carried out for claims 31-38.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/09440

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
FR 2739560	A	11-04-1997	AU 7303596 A		30-04-1997
			WO 9713524 A		17-04-1997
EP 614978	A	14-09-1994	FR 2702481 A		16-09-1994
			CA 2118558 A		10-09-1994
			JP 7149800 A		13-06-1995
US 5623053	A	22-04-1997	NONE		
WO 9640199	A	19-12-1996	AU 6386996 A		30-12-1996
			CA 2223402 A		19-12-1996
			EP 0831875 A		01-04-1998
			JP 11507824 T		13-07-1999
			US 5858981 A		12-01-1999
WO 9509002	A	06-04-1995	AU 701919 B		11-02-1999
			AU 7924494 A		18-04-1995
			CA 2172392 A		06-04-1995
			EP 0723455 A		31-07-1996
			JP 9506072 T		17-06-1997
			US 5858981 A		12-01-1999
EP 791653	A	27-08-1997	AU 3740089 A		12-12-1989
			EP 0343950 A		29-11-1989
			EP 0449828 A		09-10-1991
			JP 2657221 B		24-09-1997
			JP 3501484 T		04-04-1991
			KR 149012 B		17-08-1998
			WO 8911490 A		30-11-1989
			US 5976831 A		02-11-1999
			US 5767077 A		16-06-1998
			CA 2000355 A		19-07-1990
EP 321842	A	28-06-1989	EP 0321601 A		28-06-1989
			EP 0324879 A		26-07-1989
			AU 2738088 A		13-07-1989
			DK 712488 A		23-06-1989
			FI 885873 A		23-06-1989
			HU 50498 A		28-02-1990
			JP 2000443 A		05-01-1990
			NO 885687 A		23-06-1989
			PT 89299 A		29-12-1989
			ZA 8809506 A		29-08-1990
EP 259615	A	16-03-1988	EP 0257114 A		02-03-1988
			EP 0286700 A		19-10-1988
			AU 7716587 A		19-05-1988
			DD 263538 C		27-09-1989
			DK 434287 A		22-02-1988
			EP 0258489 A		09-03-1988
			EP 0258492 A		09-03-1988
			FI 873614 A		22-02-1988
			HU 44620 A		28-03-1988
			JP 1027496 A		30-01-1989
			NO 873512 A		22-02-1988
			NZ 221501 A		28-08-1990
			PT 85560 A, B		01-09-1987
			SU 1727533 A		15-04-1992

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/09440

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 259615 A		ZA 8706168 A	26-04-1989
WO 9905271 A	04-02-1999	AU 8455598 A	16-02-1999